

p-Block Elements (Part-2)

Group 15(Nitrogen Family)

15th group or VA group of the extended form of the periodic table consists of 5 elements.

${}^{14}_7\text{N}$ -nitrogen, ${}^{31}_{15}\text{P}$ -Phosphorus, ${}^{75}_{33}\text{As}$ -arsenic,

${}^{122}_{51}\text{Sb}$ -antimony and ${}^{209}_{83}\text{Bi}$ -bismuth. This group of five elements constitutes a family. Collectively these are known as **pnictogens** and their compounds as **pnictides**. (Greek, pniomigs meaning suffocation).

		Group		
		14 (IV A)	15 (V A)	16 (VI A)
Period	2	C (6)	N (7)	O (8)
	3	Si (14)	P (15)	S (16)
	4	Ge (32)	As (33)	Se (34)
	5	Sn (50)	Sb (51)	Te (52)
	6	Pb (82)	Bi (83)	Po (84)

Occurrence

1. Nitrogen is most abundant in nature both in free and in the combined state.
2. Air is the most abundant source of free nitrogen.
3. It forms 75% by mass and 78% by volume of the air.
4. In combined state, it is found as nitrates such as chile salt petre (NaNO_3), Indian salt petre (KNO_3) and ammonium compounds.

Concept Ladder



Nitrogen is much less reactive than phosphorous, that's why it has greater tendency for catenation than Nitrogen.

Rack your Brain



Which element shows anomalous behaviour among group 15 and why?

Previous Year's Questions



Nitrogen is relatively inactive element because

[AIPMT]

- (1) its atom has a stable electronic configuration
- (2) it has low atomic radius
- (3) its electronegativity is fairly high
- (4) Bond dissociation energy of its molecule is fairly high.

- Nitrogen is found in the form of proteins and amino acids in living organisms.
- Phosphorus occurs in minerals e.g. fluorapatite $\text{Ca}_3[\text{PO}_4]_2 \cdot \text{CaF}_2$ which is the main constituent of phosphate rocks.
- Ar, Sb and Bi are commonly associated with sulphide minerals, ores are Bi_2O_3 bismuth oxide and bismuth sulphide Bi_2S_3 .

Rack your Brain



What is the percentage of Nitrogen in air?

Electronic configuration :

Element	At. Nu.		Electronic Configuration	Brief representation of electronic configuration
Nitrogen (N)	7	2, 5	$1s^2 2s^2 2p^3$	$[\text{He}] 2s^2 2p^3$
Phosphorus (P)	15	2,8,5	$1s^2 2s^2 2p^6 3s^2 3p^3$	$[\text{Ne}] 3s^2 3p^3$
Arsenic (As)	33	2, 8, 18, 5	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^3$	$[\text{Ar}] 3d^{10} 4s^2 4p^3$
Antimony (Sb)	51	2, 8, 18, 18, 5	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^3 4d^{10} 5s^2 5p^3$	$[\text{Kr}] 4d^{10} 5s^2 5p^3$
Bismuth (Bi)	83	2, 8, 18, 32, 18, 5	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^3 4d^{10} 4f^{14} 5s^2 5p^3 5d^{10} 6s^2 6p^3$	$[\text{Xe}] 4f^{14} 5d^{10} 6s^2 6p^3$

Trends in physical and chemical properties :

Physical property

- Electronic configuration** : Valence shell electronic configuration of 15th group elements is $ns^2 np^3$, where $n = 2$ to 6.
- Atomic radii** : It increases as we move down the group. However, there is small increase in covalent radii from As to Bi, this is because of poor shielding of the valence electrons by the d- and f-electrons present in the inner shell.
- Ionization enthalpy** : Ionization enthalpy of these elements is much higher than the corresponding elements of group 14 due to increased nuclear charge and stable exactly half-filled electronic configuration.

Previous Year's Questions



Number of electrons shared in the formation of nitrogen molecule is

[AIPMT]

- (1) 6 (2) 10
(3) 2 (4) 8

- Electronegativity** : It decreases due to a gradual increase in the atomic radii of the elements.
- Melting and boiling points** : M.P and B.P. of group 15 elements first increases from N to As due to increase in their atomic size and then decreases from Sb and Bi due to their tendency to form three covalent bonds instead of five due to inert pair effect.
- Oxidation state** : These elements can show negative as well as positive oxidation states. Going down the group, the stability of +3 oxidation state increases while that of +5 oxidation state decreases due to inert pair effect.
- Non-metallic and metallic character** : Down the group, metallic character increases.

Concept Ladder



Nitrogen have higher ionisation enthalpy as compare to oxygen and carbon because of its half filled stable configuration.

Some properties of group 15 elements :

Sr.	Property	N	P	As	Sb	Bi
1	Atomic number	7	15	33	51	83
2	Valence shell configuration	$[\text{He}]^2 2s^2 2p^3$	$[\text{Ne}]^{10} 3s^2 3p^3$	$[\text{Ar}]^{18} 4s^2 4p^3$	$[\text{Kr}]^{36} 4d^{10} 5s^2 5p^3$	$[\text{Xe}]^{54} 4f^{14} 5d^{10} 6s^2 6p^3$
3	Atomic mass g/mol^{-1}	14.01	30.97	74.92	121.75	209.00
4	Covalent radius (Pm)	70	110	120	140	150
5	Ionic radius (Pm)	171 (N^{3-})	212 (P^{3-})	222 (As^{3-})	76 (Sb^{3-})	108 (Bi^{3-})
6	Electronegativity	3.00	2.10	2.20	1.82	1.67
7	Ionisation enthalpy $\Delta H / \text{kJ mol}^{-1}$	I \rightarrow 1402	1012	947	834	703
		II \rightarrow 2856	1903	1798	1595	1610
		III \rightarrow 4577	2910	2736	2443	2466
8	Density (g/cm^3)	0.879 at 63 K	1.823	5.778 (Grey form)	6.580	9.808
9	Melting point ($^{\circ}\text{C}$)	- 210	44 (White P)	614 Sublime	631	631
10	Boiling point ($^{\circ}\text{C}$)	- 196	280	610	1587	1564

Allotropy:

All the elements except bismuth show allotropy. Phosphorus exists in three allotropic forms such as white or yellow, red or violet and black phosphorus.

Rack your Brain



How Phosphorus is categorised into its white, red and black allotropic form?


Property	White phosphorus	Red phosphorus	Black phosphorus
Colour	White, but turns yellow on exposure	Dark red	Black
State	Waxy solid, can be cut with knife	Brittle powder	Crystalline with greasy touch
Smell	Garlic smell	Odourless	—
Density	1.849 cm ⁻³	2.19 cm ⁻³	2.69 g cm ⁻³
Ignition Temperature	307 K	543 K	673 K

Chemical properties :

1. Action of air :



2. Action of oxidising acids :



3. Hot concentrated nitric acid and sulphuric acid has no reaction with nitrogen.



Concept Ladder



- Dinitrogen combines with dioxygen only at very high temperature (2000 K) to form nitric oxide.
- Dinitrogen is a colourless, odourless, tasteless and non-toxic gas.

4. Action of alkalis :



Salt



Salt

Alkalies have no reaction with nitrogen.

5. Action of metals :



Fulfillment of Octet

The elements of group 15 have $ns^2 np^3$ as their valence shell electronic configuration.

They can complete their octets in two different ways:

(a) Electron Transfer:

The atoms of the elements of this group may accept three electrons from more metallic elements to form triply charged negative ions such as nitride, N^{3-} ion and phosphide, P^{3-} ion and thereby attain noble gas configuration. Only small atoms can form highly charged negative ions because of their greater electronegativities. The other members of nitrogen family show little tendency to form triply charged negative ions and this tendency decreases down the group because of increase in size and decrease in electronegativity. These group elements also exhibit oxidation states ranging from -3 to +5.

Concept Ladder



Oxides of nitrogen have open chain structure, while those of phosphorous have closed chain or cage structures.

Rack your Brain



Why NO_2 dimerises to form N_2O_4 ?

Previous Year's Questions



Which of the following oxides is most acidic?

[AIPMT]

(1) As_2O_5

(2) P_2O_5

(3) N_2O_5

(4) Sb_2O_5

(b) Electron Sharing:

Since the atoms of these elements contain three unpaired p-electrons so these can pair with unpaired p-electrons in another atom or atoms to form three covalent bonds, e.g., NH_3 , PH_3 , AsH_3 , SbH_3 , BiH_3 .

Compounds formed by Group 15 Elements

Hydrides :

Stability order : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$

Basicity : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$

H-E-H bond angle : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3$


Boiling point : $\text{SbH}_3 > \text{NH}_3 > \text{AsH}_3 > \text{PH}_3$

Reducing character : $\text{BiH}_3 > \text{SbH}_3 > \text{AsH}_3 > \text{PH}_3 > \text{NH}_3$

The bond angle in PH_3 , AsH_3 and SbH_3 is close to 92° which suggests that orbitals used for bonding are close to pure p-orbitals.

Oxides :

Oxides with higher oxidation state are more acidic :



Moreover, acidity decreases down the group :



Concept Ladder



Tendency to form pentahalides decreases down in group 15 elements.

Rack your Brain



Nitrogen does not form pentahalides, why phosphorus does so?

Q.1 What is fuming nitric acid? What colour is it have?

A.1 Concentrated nitric acid in which oxides of nitrogen (i.e., NO , NO_2 , N_2O_3 , etc.) are dissolved is called fuming nitric acid. Its colour is yellow.

Chemical Properties

(a) Ammonia (NH₃)


Ammonia is a colorless gas having characteristic pungent smell. It is quite poisonous and brings tears in eyes. NH₃ is highly soluble in water due to extensive hydrogen bonding.

Preparation:

1. In laboratory:


NH₃ produced is dried with quick lime CaO. Note that it cannot be dried with H₂SO₄, CaCl₂ or P₄O₁₀ as they react with the gas instead of drying it.

2. Haber - Bosch Process :




$$\Delta H^\ominus = -46.1 \text{ kJ / mol}$$

3. NH₃ is prepared by the Cyanamide process :



4. Serpeck's Process (metallurgy of aluminium) :



Chemical Properties:

1. Ammonia is a non-supporter of combustion but burns in oxygen with a pale-yellow flame if continuous heat is supplied.




Q.2

What happens when a mixture of ammonium sulphate and sodium nitrate is heated. Write the equation.

A.2

Double displacement reaction first occurs to form ammonium nitrate which then decomposes to yield laughing gas (N₂O).



Concept Ladder



Haber process also known as artificial nitrogen fixation, is the main industrial procedure for the production of ammonia.







Previous Year's Questions

Pure nitrogen is prepared in the laboratory by heating a mixture of
[AIPMT]


- (1) NH₄OH + NaCl
- (2) NH₄NO₃ + NaCl
- (3) NH₄Cl + NaOH
- (4) NH₄Cl + NaNO₂




2. Formation of complex ions used in qualitative analysis:



3. Reaction with Nessler's Reagent is used as a test to detect ammonia:




4. Reaction with chlorine:



(b) Ammonium Salt

All ammonium salts are very soluble in water. They are usually slightly acidic and decompose readily on heating producing NH_3 , N_2 or N_2O .

- If the anion (B^-) is not particularly oxidising (eg. Cl^- , CO_3^{2-} or SO_4^{2-}) the ammonia is evolved.



Concept Ladder



Due to H-bonding NH_3 exists in both solid and liquid states and has higher melting and boiling points than expected on the basis of its molecular mass.






Previous Year's Questions

Which one has the lowest boiling point?

[AIPMT]

- (1) NH_3 (2) PH_3
(3) AsH_3 (4) SbH_3

- If the anion is more oxidising (e.g. NO_2^- , NO_3^- , ClO_4^- , $\text{Cr}_2\text{O}_7^{2-}$) than NH_4^+ is oxidized to N_2 or N_2O .



(c) Oxides of Nitrogen:





1. Nitrous oxides N_2O [Laughing gas]:

It is colourless, non-combustible gas with a sweetish odour and taste. It is a neutral oxide and is diamagnetic.




Preparation:



(i) It is prepared by careful thermal decomposition of molten ammonium nitrate at about 280°C



Chemical Properties:



(b) Reduction of N_2O to N_2 :



Concept Ladder



The bond angle ($\text{O}-\text{N}-\text{O}$) are not of the same value in NO_2^- and NO_2^+ .

Rack your Brain



Write the name and formula of neutral oxides of nitrogen.

Previous Year's Questions




Which of the following oxides of nitrogen is paramagnetic?

[AIPMT]

- (1) NO_2 (2) N_2O_3
 (3) N_2O (4) N_2O_5



(c) Oxidation:




Uses:

It is used as an anaesthetic by dentists and a propellant for whipped ice-cream.

2. Nitric Oxide (NO):

It is a colourless, neutral gas which is paramagnetic due to the presence of odd electrons. It is sparingly soluble in water.



Preparation


(a) NO is prepared in the laboratory by the reduction of dilute HNO_3 with Cu.



(b) Commercial Method




(In Birkeland and Eyde Process)



(Catalytic Oxidation of Ammonia)

Chemical Properties:

(i) It is sparingly soluble in water and its solid-state forms diamagnetic dimer



(ii) $\text{NO} + 1/2\text{O}_2 \rightarrow \text{NO}_2$ (brown fumes), thus it is not possible to detect its smell.

[This reaction is also used to detect O_2]

Concept Ladder



NO is paramagnetic in its gaseous state but diamagnetic in its liquid or solid state.

Rack your Brain




Why in the structure of HNO_3 molecule, the N—O bond (121 pm) is shorter than N—OH (140 pm).


(iii) Action of heat:



(iv) Reduction (to N_2):



(v) Oxidation/Reduction properties





3. Dinitrogen Trioxide N_2O_3 :

It exists only in solid state at low temperature which is pale blue in colour. N_2O_3 is an acidic anhydride of nitrous acid (HNO_2). It is diamagnetic.



Preparation:

(a) It can be made by condensing equimolar amounts of NO and NO_2 together at 253 K.



Concept Ladder



N_2O_3 i.e. Nitrogen trioxide, is a highly toxic compound and can be partially dissociated into NO & NO_2 .

Rack your Brain



Why N_2O is diamagnetic in nature?

Previous Year's Questions




Which of the following is a nitric acid anhydride?

[AIPMT]


- (1) NO (2) NO_2
(3) N_2O_5 (4) N_2O_3



Chemical Properties:




(ii) It being the anhydride of nitrous acid forms nitrites with alkalis.




(iii) It forms brown coloured $\text{FeSO}_4 \cdot \text{NO}$ with FeSO_4 . This is used as a test for nitrates called Brown Ring Test.

4. Nitrogen Dioxide NO_2 :

It's a reddish-brown gas and exists at the room temperature. Being an odd electron molecule, it is paramagnetic and dimerises into N_2O_4 at low temperature. It is acidic in nature.



Structure:



Concept Ladder



NO_2 is coloured but its dimer form N_2O_4 is colourless in nature.

Rack your Brain




Why NO_2 is called as mixed anhydride of nitrous acid?

Q.3 Why does NO_2 dimerize?


A.3 NO_2 is an odd electron ($7 + 2 \times 8 = 23$) molecule. In the valence shell, N has seven electrons and hence is less stable. To become more stable by acquiring inert gas configuration having 8 electrons in the valence shell, it undergoes dimerization to form N_2O_4 .

Separation:


(a) In the laboratory it is prepared by heating dried lead nitrate:



(b) It is produced on a large scale by oxidizing NO in the Ostwald's Process for the manufacture of nitric acid.



Chemical Properties:





Thus, NO_2 is called as mixed anhydride of nitrous acid and nitric acid. The reaction of NO_2 with aqueous alkalis is similar to that of a mixture of HNO_2 and HNO_3 .



(ii) Oxidation:




(iii) Reduction:



5. Dinitrogen Pentoxide N_2O_5 :

N_2O_5 is a colourless crystalline solid which sublimates readily and is acidic.



Concept Ladder



NO_2 is an intermediate in the industrial synthesis of nitric acid, it is paramagnetic, bent in shape and used in fertilizers.

Previous Year's Questions




When copper is heated with conc. HNO_3 it produces

[NEET-2016]



- (1) $\text{Cu}(\text{NO}_3)_2$, NO and NO_2
- (2) $\text{Cu}(\text{NO}_3)_2$ and N_2O
- (3) $\text{Cu}(\text{NO}_3)_2$ and NO_2
- (4) $\text{Cu}(\text{NO}_3)_2$ and NO

The solid form consists of $(\text{NO}_2^+ + \text{NO}_3^-)$ which is ionic and called as nitronium nitrate while in solution it is covalent.

Preparation:



Thus N_2O_5 is the anhydride nitric acid.



Chemical Properties


(i) N_2O_5 dissolves in H_2O with a hissing sound.



Thus it is called as nitric anhydride and neutralisation reactions similar to HNO_3




(ii) Since $\text{N}_2\text{O}_5 \xrightarrow{\Delta} \text{NO}_2 + \text{NO} + \text{O}_2$ it is a very powerful oxidising agent. It destroys all organic substances.



(d) Oxoacids of Nitrogen:

1. Nitrous acid HNO_2 :

It is unstable except in dil. aq. solution. It is a tautomeric mixture of two forms:



Concept Ladder



N_2O_5 i.e. Nitric anhydride is an ore of binary nitrogen oxides, a family of compounds that only contain nitrogen and oxygen.

Rack your Brain



Find the covalency of N in N_2O_5 ?

Previous Year's Questions




Urea reacts with water to form A which will decompose to form B. B when passed through $\text{Cu}^{2+}_{(aq)}$, deep blue colour solution C is formed. What is the formula of C from the following?

[NEET-2020]


- (1) CuSO_4 (2) $[\text{Cu}(\text{NH}_3)_4]^{2+}$
 (3) $\text{Cu}(\text{OH})_2$ (4) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$

Preparation:

(a) It is made by acidifying a solution of a nitrite





(b) Dissolving nitrous anhydride (N_2O_3) in H_2O of course forms HNO_2 :



Oxidising Properties:


NO_2^- ion is a weak oxidising agent and gets reduced to NO which forms a red complex with haemoglobin and improves the look of meat.




However, it is oxidised by KMnO_4 , Cl_2 forming NO_3^- (but to NO_3 by H_2SO_4):




Reaction with Amines :



Secondary and tertiary aliphatic amines form nitrosamines with nitrites:



Nitrites are used to make diazo compounds used in azo dyes:



Concept Ladder



The N—O bond in NO_2^- is shorter than N—O bond in NO_3^- .

Rack your Brain



Draw the structure of N_2O_5^- .




Other Reactions:

On standing HNO_2 decomposes into NO and NO_2 :



While on boiling it disproportionates:



2. Nitric acid HNO_3 :

Also called as Aqua fortis, it is colourless fuming liquid with a pungent smell. Yellow colour occurs on standing due to decomposition into NO_2 .




Preparation:

1. In laboratory





HNO_3 vapours are then condensed in a water-cooled receiver.

2. Birkeland and Eyde Process



Q.4 How is nitrogen prepared from ammonia?

A.4 Nitrogen is prepared by ammonia according to the following equations :




Concept Ladder




N_2O_5 is a rare example of a compound that adopts two structures depending on the conditions. The solid form is a salt in which nitronium nitrate, consisting of separate nitronium cations $[\text{NO}_2]^+$ and nitrate anions $[\text{NO}_3]^-$.

3. Ostwald Process




Due to the formation of azeotrope, HNO_3 cannot be further concentrated by distillation and other methods are used like dehydrating using sulphuric acid vapours.



Oxidising Properties:

NO_3^- is a very powerful oxidising agent in acidic solution. Cu, Ag which are insoluble in HCl dissolve in HNO_3 . Concentrated HNO_3 forms NO_2 with a reducing agent although. Fe, CO, Al, Ni, Cr and B are rendered passive by concentrated HNO_3 due to the formation of a protective oxide layer (like Fe_3O_4 with iron). Dilute HNO_3 forms NO with a poor reducing agent (Cu, Hg); N_2O with a good reducing agent in hot conditions and NH_4NO_3 with a good reducing agent in cold conditions.



Previous Year's Questions



Zn gives H_2 gas with H_2SO_4 and HCl but not with HNO_3 because

[AIPMT]

- (1) Zn act as oxidising agent when react with HNO_3
- (2) HNO_3 is weaker acid than H_2SO_4 and HCl
- (3) in electrochemical series Zn is above hydrogen
- (4) NO_3^- is reduced in preference to hydronium ion.







Rack your Brain



How conc. HNO_3 turns yellow on exposure to sunlight.



Some reactions with non-metals are:



- Only Mg and Mn react with 2% dil. HNO_3
 HNO_3 (dil. 2%) + Mg (or Mn) \longrightarrow $\text{Mg}(\text{NO}_3)_2$ + $\text{H}_2 \uparrow$
- Solution of 75% HCl and 25% HNO_3 is called Aqua Regia which contains NO^+ Cl^- ions. Aqua Regia is capable of dissolving Gold and Platinum by forming soluble H_3AuCl_4 and H_2PtCl_6 respectively.

Other Oxoacids:

Other Oxoacids	
HOONO	Pernitrous acid
H_2NO_2	Hydronitrous acid
HNO_4	Pernitric acid
$\text{H}_2\text{N}_2\text{O}_2$	Hyponitrous acid

Concept Ladder



Aqueous HNO_3 can be concentrated by distillation upto 68% by mass. Further concentration to 98% can be achieved by dehydration with conc. H_2SO_4 .

Rack your Brain



Draw structure of compounds H_2NO_2 and $\text{H}_2\text{N}_2\text{O}_2$. Is there any possibility of having double bond between N and O?

Q.5 What is the product formed by the following reaction?




A.5 P_4O_{10} being a strong dehydrating agent, eliminates a molecule of H_2O from two molecules of HNO_3 giving dinitrogen pentoxide (N_2O_5).



Phosphorous and its Compounds :

Allotropy :

It exists in many allotropic forms, the important ones being white, red and black phosphorous.



Rack your Brain



Why Phosphorous only exist in cage or close chain structure?

White phosphorus	Red phosphorus
It is a white, waxy and highly toxic solid. It glows in dark (Chemiluminescence/Phosphorescence)	It is non poisonous and does not show phosphorescence
It is unstable due to angular strain and spontaneously reacts with O ₂ and catches fire. $\text{P}_4 + 5\text{O}_2 \longrightarrow \text{P}_4\text{O}_{10}$ Thus it is stored in water in which it is insoluble (white phosphorous is soluble in CS ₂)	It is stable in air and not stored in water. It is insoluble in organic solvents.
 Tetrahedral	

Only white phosphorous reacts with caustic alkalis to undergoes a disproportion reaction.




Uses :

Red phosphorus is used in Match - Industry and white phosphorus as a rat poison.

Preparation:




Phosphorous is obtained by the reduction of calcium phosphate with C in an electric furnace. Sand (SiO_2) is added to remove the calcium as a fluid slag.



Overall reaction :



Oxides of Phosphorus

Phosphorous Trioxide [$\text{P}_2\text{O}_3/\text{P}_4\text{O}_6$]	Phosphorous Pentoxide [$\text{P}_2\text{O}_5/\text{P}_4\text{O}_{10}$]
It is also called Phosphorus oxide or phosphorous anhydride. It is a soft white solid.	It is called phosphoric oxide or Phosphoric anhydride.
Phosphorous trioxide is dimeric and written as P_4O_6 	
Structure of phosphorus trioxide P_4O_6	The shorter $\text{P} \rightarrow \text{O}$ coordinate bond is in fact a 'double bond' different from the usual double bond. A full p orbital on O overlap side ways with an empty d orbital on p atom forming $\text{p}\pi\text{-d}\pi$ back bonding or a dative bond. 

Previous Year's Questions




Which of the following phosphorus is the most reactive?

[AIPMT]

- (1) Scarlet phosphorus
- (2) White phosphorus
- (3) Red phosphorus
- (4) Violet phosphorus

Other oxides like P_4O_8 and P_4O_9 are intermediate between P_4O_6 and P_4O_{10} and form a mixture of phosphoric acid P(+V) and phosphorous acid P(+III) on hydrolysis.



Oxo Acids of Phosphorous


All oxo acids of phosphorous can be categorized into two main acid series namely, phosphorus acid series and phosphoric acid series.

The Phosphorus acid series:

The series contain P(+3) and are generally reducing agents due to the presence of P – H bonds which are reducing (as they break down easily to loose H^+).

(I) Ortho phosphorous acid H_3PO_3 :


It is a dibasic acid formed by hydrolysis of P_4O_6 .



It forms phosphites on hydrolysis which are very strong reducing agents in basic solutions.



On heating orthophosphorous acid disproportionates :



Concept Ladder



Phosphinic acid behaves as a mono protic acid where as Ortho Phosphorus acid is diprotic acid.

Rack your Brain



Find the product formed when PCl_5 reacts with heavy water.

Previous Year's Questions



Which is the correct statement for the given acids?


[NEET-2016]

- (1) Phosphinic acid is a mono protic acid while phosphonic acid is a diprotic acid.
- (2) Phosphinic acid is a diprotic acid while phosphonic acid is a monoprotic acid.
- (3) Both are diprotic acids
- (4) Both are triprotic acids.




(II) Hypophosphorous Acid H_3PO_2 :


Hypophosphorous acid is prepared by alkaline hydrolysis of (white) phosphorous.




Structure of P_4



H_3PO_2 is a monobasic acid and a very strong reducing agent.



(III) Pyrophosphorous Acid $\text{H}_4\text{P}_2\text{O}_5$:



While reaction with PCl_5 is



Concept Ladder



$\text{H}_4\text{P}_2\text{O}_5$ is dibasic as metals can replace the hydrogen in the pair of attached hydroxide ions.

Q.6 H_3PO_3 undergoes disproportionation reaction but H_3PO_4 does not?

A.6 The maximum and minimum oxidation states of P are -3 and $+5$. But the oxidation state of P in H_3PO_3 is $+3$. therefore, it can increase its oxidation to $+5$ in H_3PO_4 and decrease its oxidation state to -3 in PH_3 .

Thus, H_3PO_3 shows disproportionation reaction : $4 \text{H}_3\overset{+3}{\text{P}}\text{O}_3 \longrightarrow 3\text{H}_3\overset{+5}{\text{P}}\text{O}_4 + \text{PH}_3$

In contrast, the oxidation state of P in H_3PO_4 is $+5$, therefore, it cannot increase its oxidation state beyond $+5$ and hence it does not show disproportionation reaction.


The Phosphoric Acid Series:

They usually contain P(+V) and have oxidising properties. The simplest phosphoric acid is orthophosphoric acid.

(I) Orthophosphoric acid H_3PO_4 :

Preparation:


It is formed by hydrolysis of phosphorous pentoxide (P_2O_5 or P_4O_{10}). The scheme for its hydrolysis is:



In laboratory



Impure H_3PO_4 is prepared in large amounts by 'Wet Process'.



$CaSO_4$ is hydrated to gypsum $CaSO_4 \cdot 2H_2O$ and filtered off. H_3PO_4 thus produced is used to make fertilizers.

Properties:

1. H_3PO_4 is hydrogen bonded in aqueous solution and thus the 'concentrated acid' is syrupy and viscous.

Rack your Brain



How many P—O—P bonds are there in $H_4P_2O_7$?

Previous Year's Questions




P_2O_5 is heated with water to give

[AIPMT]

- (1) hypophosphorous acid
- (2) phosphorous acid
- (3) hypophosphoric acid
- (4) orthophosphoric acid




2. It forms meta phosphates on heating:




Sodium metaphosphate is also termed Graham's salt or Calgon (Commercial name). It is soluble in H_2O and precipitates Pb^{2+} , Ag^+ but not Ca^{2+} , Mg^{2+} . Thus, it is used for softening water.

3. It forms 3 series of salts:



Dihydrogen phosphates are slightly acidic in water.



Monohydrogen phosphates are slightly basic in water.




Normal phosphates are appreciably basic in water.

4. Phosphates are estimated quantitatively by adding a solution containing NH_4OH solution of the phosphate and Mg^{2+} . Magnesium ammonium phosphate precipitated is ignited and weighed as $\text{Mg}_2\text{P}_2\text{O}_7$.



(II) Hypophosphoric acid $\text{H}_4\text{P}_2\text{O}_6$:

$\text{H}_4\text{P}_2\text{O}_6$ contains P(+4) and is prepared by hydrolysis and oxidation of red phosphorous by NaOCl .



Concept Ladder




- H_3PO_4 i.e. phosphoric acid has its pure compound in colourless solid form.
- When all three H^+ ions removed, orthophosphate ion PO_4^{3-} is form, which is also commonly known as phosphate.

Rack your Brain




Name the compound formed as esters by orthophoric acid.

On hydrolysis $\text{H}_4\text{P}_2\text{O}_6$ forms both H_3PO_3 and H_3PO_4



Phosphine PH_3


PH_3 is a colourless, extremely toxic gas which smells of rotten fish. It is highly reactive and not soluble in H_2O . Its aqueous solutions are neutral.



The bond angle in $\text{H}-\text{P}-\text{H}$ is 93.5° which suggests the presence of almost pure p orbitals.

Preparation:

PH_3 can be formed by hydrolysing metal phosphides or hydrolysing white phosphorous in basic media.



Another method also produces small amounts of highly inflammable P_2H_4 . Therefore, PH_3 is removed in the following manner:



Properties :

Q.7 PH_3 has lower boiling point than NH_3 . Why?

A.7 The electronegativity of N (3.0) is much higher than that of P (2.1). Therefore, NH_3 undergoes extensive intermolecular H-bonding and hence it exists as an associated molecule. To break these H-bonds, a large amount of energy is needed. On the other hand, PH_3 does not undergo H-bonding and thus exists as discrete molecules. Therefore, the boiling point of PH_3 is much lower than that of NH_3 .

Concept Ladder




A French chemist Philippe once generated a regular supply of heat over Phosphorous (P_4) with an aqueous solution of K_2CO_3 , results in colourless gaseous substance called phosphine (due to its rotting fish odour).

Rack your Brain



What is the bond angle between $\text{H}-\text{P}-\text{H}$ in phosphine?

1. Phosphine is stable in air but catches fire at 150°C.





2. It explodes in contact with traces of oxidising agents like HNO_3 , Cl_2 and Br_2 .




3. Solution of PH_3 in H_2O decomposes in presence of light giving red P and H_2 .

4. Formation of metallic phosphides:



5. PH_3 is weakly basic and forms phosphonium salts with anhydrous acids while NH_3 readily forms NH_4X in aqueous solutions of the acids.





Uses:

It is used for making 'Holme's signal'. CaC_2 and Ca_3P_2 are contained in containers. Containers are pierced and thrown in sea when gases evolved, burn and serve as a signal.

Halides:

1. PCl_5 is a yellowish white powder while PCl_3 is a colourless liquid and is widely used in organic chemistry.



Concept Ladder



PH_3 possesses the characteristics to cause high environmental damages, it acts as a Lewis base when its lone pair electron is donated in reaction with hydrogen iodide.

Rack your Brain



Why PCl_4^+ can exist but PCl_2^- cannot?


Previous Year's Questions




A compound 'X' upon reaction with H_2O produces a colourless gas 'Y' with rotten fish smell. Gas 'Y' is absorbed in a solution of CuSO_4 to give Cu_3P_2 as one of the products. Predict the compound 'X'.

[NEET-2019]


- (1) Ca_3P_2 (2) NH_4Cl
 (3) As_2O_3 (4) $\text{Ca}_3(\text{PO}_4)_2$



2. PCl_3 fumes in moisture.



3. In solid state it is ionic therefore conducts electricity, while in gaseous and liquid state PCl_5 is trigonal bipyramidal. While in solid state it is ionic and hence conducts electricity



4. PF_5 forms an unusual trigonal bipyramid where equatorial and axial bonds interchange their positions in short time. This is called pseudo rotation.


Fertilizers:

They are the compounds of N, P and K which are soluble in H_2O and easily assimilated by plants without causes any damage to them.


I. Nitrogenous Fertilizers:

(i) Ammonium sulphate $((NH_4)_2SO_4)$:

Ammonium sulphate is made by passing NH_3 and CO_2 gases into a slurry of $CaSO_4$ in water:



In soil, the action is:



Concept Ladder



The main use of ammonium sulphate is as a fertilizer for alkaline soils. Ammonium ion forms a small amount of acid, which lowers pH balance of the soil.

Rack your Brain



Why calcium ammonium nitrate does not form soil acid where as ammonium sulphate do?

Group 16 (Oxygen family)

Introduction :

The elements Oxygen (O), Sulphur (S), Selenium (Se), Tellurium (Te) and Polonium (Po) constitute group 16 of the periodic table.

Group 16 elements are also known as Chalcogens, means ore forming, because many metals are found as oxides and sulphides and a few as selenides and tellurides.

Occurrence :

1. Oxygen is the most abundant element and is found both in free as well as in combined state.
2. Oxygen makes up 46.6% by mass of the earth's crust and dry air contains 21.0% oxygen by volume.
3. Sulphur occurs less abundantly and constitutes 0.034% by mass of the earth's crust.
4. Sulphur occurs mainly in combined form of sulphates and sulphides.
5. Onion, garlic, mustard, eggs, proteins hair and wool, the organic materials also contain sulphur.
6. Polonium is radioactive in nature.

Electronic configuration :

Element	symbol	Atomic Number	Atomic Electronic Configuration	Brief representation of electronic configuration
Oxygen	O	8	$1s^2 2s^2 2p^4$	$[\text{He}] 2s^2 2p^4$
Sulphur	S	16	$1s^2 2s^2 2p^6 3s^2 3p^4$	$[\text{Ne}] 3s^2 3p^4$
Selenium	Se	34	$1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4$	$[\text{Ar}] 3d^{10} 4s^2 4p^4$
Tellurium	Te	52	$1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 5s^2 5p^4$	$[\text{Kr}] 4d^{10} 5s^2 5p^4$
Polonium	Po	84	$1s^2 2s^2 2p^6 3s^2 3p^4 3d^{10} 4s^2 4p^4 4d^{10} 4f^{14} 5s^2 5p^4 5d^{10} 6s^2 6p^4$	$[\text{Xe}] 5d^{10} 6s^2 6p^4$

Concept Ladder



Sulphur has greater tendency for catenation than oxygen.

Rack your Brain



Arrange the ions of group 16 elements according to their atomic size.

Trends in physical and chemical properties :


Physical properties :

1. Physical state :

Oxygen is gas while all others are solids.

2. Molecular structure :

Oxygen molecule is diatomic while the molecules of other element are more complex. Sulphur, selenium and tellurium exists as staggered 8-atom rings.



Crown shaped puckered ring of S_8 .

3. Atomic and ionic radii :

It increases on moving down the group due to increase in the number of electron shells.

4. Ionization enthalpy :

The ionization energies of the elements of oxygen family are less than those of nitrogen family. This is due to the reason that pnicogens has exactly half-filled orbitals and thus configuration is stable. As we move from oxygen to polonium, the ionization energy decreases due to increase in atomic size.

5. Melting and boiling point :

M.P. and B.P. increases with the increases in atomic numbers because of the increase in magnitude of Vander waals forces.

6. Electronegativity :

Elements of group 16 are more electronegative than corresponding elements of group 15 due to their smaller atomic size.

7. Metallic and non-metallic character :

They have very less metallic character due to high ionisation energy. Their metallic character increases down the group from oxygen to polonium because of decrease in ionization energy.

Concept Ladder



The value of electron gain enthalpy with negative sign for sulphur is higher than that of oxygen.

Rack your Brain



Name the element(s) of group 16 which is amphoteric in nature.

Previous Year's Questions



Which is the correct thermal stability order for H_2E ($E = O, S, Se, Te$ and Po)?

[NEET-2019]

- (1) $H_2Se < H_2Te < H_2Po < H_2O < H_2S$
- (2) $H_2S < H_2O < H_2Se < H_2Te < H_2Po$
- (3) $H_2O < H_2S < H_2Se < H_2Te < H_2Po$
- (4) $H_2Po < H_2Te < H_2Se < H_2S < H_2O$

8. Electron gain enthalpy :


The elements of oxygen family have high electron gain enthalpy. The value decreases from sulphur to polonium.

9. Allotropy :

All the elements of 16th group show allotropy. Oxygen exists in two non-metallic forms i.e. O₂ and O₃. Sulphur provides a very good example of an element that exhibits allotropy.

10. Catenation :

Oxygen has some but sulphur has greater tendency for catenation.



Concept Ladder



The boiling point order among H₂O, HCl and NH₃ is H₂O > NH₃ > HCl, it is because H₂O is capable of showing more stronger H-bonding than NH₃, while HCl has no H-bonding.

Some properties of group 16 elements.

Properties	O	S	Te	Se	Elements Po
Atomic No.	8	16	34	52	84
Atomic mass	16.00	32.06	77.96	127.60	210
Covalent radius (pm)	74	104	117	137	140
Ionic radius M ⁻² (pm)	140	184	198	221	230(approx)
Density at 298 K (g cm ⁻³)	1.32 (at m.p.)	2.06 (for- rhombic form)	4.19 (for hex agonal gray allotrope)	6.25	—
Melting point K	55	386 (for momo- clinic form)	490	723	527
Boiling point K	90	718	958	1263	1233
Ionization enthalpy (I) KJ mol ⁻¹	1310	1000	941	870	812
Electronegativity	3.50	2.5	2.48	2.1	2.0
Electron gain enthalpy DH / kJ mol ⁻¹	-141	-200	-195	-190	-183



Oxygen and its compounds



Preparation :

O₂ is a colourless, odourless and tasteless gas which is partially soluble in water. Liquid dioxygen is pale blue in colour and the solid is also blue. It is paramagnetic due to presence of unpaired electrons.

1. It is prepared industrially by the electrolysis of acidic or alkaline H₂O or by Claude's Process (fractional distillation of liquid air)
2. In laboratory it is prepared from thermal decomposition of



3. It can also be made by catalytic decomposition of hypochlorites:



Uses:

Most of the O₂ is used in the steel making industry. Some of it is used in oxy-acetylene welding and as an oxidant in rockets. It is an essential constituent of life.



Concept Ladder



Oxygen does not show oxidation state of +4 and +6, which S does so.



Previous Year's Questions

Which of the following does not give oxygen on heating?

[NEET-2013]

- | | |
|---|--|
| (1) K ₂ Cr ₂ O ₇ | (2) (NH ₄) ₂ Cr ₂ O ₇ |
| (3) KClO ₃ | (4) Zn(ClO ₂) ₂ |

Q.8 Why OF₆ does not exist but SF₆ exists?

A.8 Due to absence of d-orbitals in the valence shell, oxygen cannot show an oxidation state of +6 and hence does not form OF₆. In contrast, S contains d-orbitals in the valence shell and hence can show a maximum oxidation state of +6. Since F₂ is a very strong oxidising agent, it oxidises sulphur to its maximum oxidation state of +6 and thus forms SF₆.




Preparation:

O₃ is prepared by the action of a silent electric discharge upon dioxygen in an ozoniser.


Bond order = 1.5 for O – O bonds.

Chemical Properties:



- O₃ turns starch iodide paper blue.
- The amounts of O₃ in a gas mixture is determined by passing gas into KI solution buffered with a borate buffer (pH 9.2) Iodine liberated is titrated with sodium thiosulphate.





Alternatively, it is decomposed catalytically and the change in volume measured.



- Hg in the presence of ozone is oxidized to sub-oxide and starts sticking to glass and loses its meniscus. This is used as a test for ozone called 'Tailing of Mercury'.
- O₃ is an extremely powerful oxidising agent, second only to F₂.



- O₃ reduces peroxides



Uses

O₃ is used as a disinfectant for water. Its advantage over chlorine is that it avoids unpleasant smell and taste of chlorine, since any excess O₃ decomposes to O₂.



Concept Ladder



O₃ is used as a disinfectant and as a germicide. It is also used for purifying air in crowded places such as cinema halls, tunnels and underground railway station.



Rack your Brain

Why two O—O bond length in the ozone molecule are same?



Previous Year's Questions

The angular shape of ozone molecule (O₃) consists of

[AIPMT]



- 1 sigma and 1 pi bond
- 2 sigma and 1 pi bond
- 1 sigma and 2 pi bond
- 2 sigma and 2 pi bond

[c] Hydrogen Peroxide H_2O_2


Pure H_2O_2 is a colourless liquid (like water), more hydrogen bonded than water (and thus has highest boiling point)

Preparation:

1. At one time H_2O_2 was obtained by electrolysis of H_2SO_4 or $(NH_4)_2SO_4$ using high current density.





2. Now, it is produced industrially by a cyclic process.




H_2O_2 is extracted with water as 1% solution which is concentrated to 30% solution.

3. In laboratory



Chemical Properties :

1. H_2O_2 decomposes in presence of impurities like Fe^{2+} , Fe^{3+} , Ni^{2+} , Pt. to undergoes a disproportionation reaction :



Glycerol or acetanilide is added to check its decomposition.

Concept Ladder



H_2O_2 is used as an oxidizer, bleaching agent and antiseptic.

Rack your Brain



How H_2O_2 can be produced by using cyclic process?



Previous Year's Questions



The O—O—H bond angle in H_2O_2 is
[AIPMT]

- (1) 106° (2) $109^\circ 28'$
(3) 120° (4) 97°

2. In most of its reactions H_2O_2 acts as a strong oxidising agent (slow in acidic medium while fast in alkaline).




3. H_2O_2 is forced to act as a reducing agent with stronger oxidising agents and O_2 is evolved.




H_2O_2 is useful to counteract chlorine:

4. H_2O_2 is a mild bleaching agent for hair feathers due to the oxidation reaction:




5. Qualitative Analysis: It is detected qualitatively by its action on KI.




Structure:

It is proposed H_2O_2 is a tautomeric mixture of two forms:



It shows restricted rotation about O—O bond due to repulsion between two H's. The structure is same in liquid and gaseous forms, but angles and bond lengths change due to hydrogen bonding.



Structure of H_2O_2 in gas phase

Concept Ladder



H_2O_2 is a reactive oxygen species and the simplest peroxide, a compound having an oxygen – oxygen single bond structure.

Rack your Brain



Name the structure of H_2O_2 ?

Previous Year's Questions



Hydrogen peroxide molecules are

[AIPMT]

- (1) monoatomic and form X_2^{2-} ions
- (2) diatomic and form X^- ions
- (3) diatomic and form X_2^- ions
- (4) monoatomic and form X^- ions

Sulphur and its Compounds

Allotropy:

Sulphur has the maximum number of allotropic structures. The two common crystalline forms are :

(i) α -Rhombic Sulphur:

It is a yellow solid stable at room temperature, M.P. 11.8°C and specific gravity 2.06 g/cc.


(ii) β - Monoclinic Sulphur:

It is a colourless solid stable above 95.5°C, M.P. 120°C and specific gravity 1.98g/cc.

At 95.5°C (transition temperature both forms are stable). Both α and β forms contain puckered S_8 rings in crown conformation.

(iii) Engels Sulphur (Σ - sulphur):

It is unstable and has S_8 rings in chair conformation.




(iv) Plastic Sulphur (χ - sulphur):

It is elastic, dark coloured and obtained liquid sulphur into water. It can be moulded with fingers.

(v) Colloidal Sulphur:

It is prepared by passing H_2S gas into an oxidising solution like HNO_3 , FeCl_3



Extraction:

Sulphur is obtained from natural gas plants (which contain) and from metal sulphide ores (especially iron pyrites) which produce SO_2 .



Uses

Almost 90% of S produced is used to manufacture H_2SO_4 . The rest is used in vulcanizing of rubber, making fungicides, gunpowder etc.

Concept Ladder



α -sulphur is insoluble in water but dissolves to some extent in benzene, alcohol and ether.

Rack your Brain



How mono-clinic sulphure is prepared. Draw its structure?

Concept Ladder





α -sulphur is stable below 369 K and transform into β -sulphur above this temperature.

[a] Hydrogen sulphide H_2S (Sulphuretted Hydrogen)

It is a colourless, poisonous gas, soluble in H_2O and smell of rotten eggs.


Preparation:

It is easier to make H_2S by the action of mineral acids on metal sulphides.



Chemical Properties:



1. It burns with blue flame.



A saturated solution of H_2S is slowly oxidized by air and Sulphur is deposited.



2. It is a very weak dibasic acid.




The two salts $NaHS$, Na_2S are basic and soluble in water.

3. $H_2S + \underset{\text{dil. solution}}{NH_3} \rightarrow (NH_4)HS$ [not $(NH_4)_2S$]

Yellow $(NH_4)_2S$ is a mixture of ammonium polysulphide and is made by dissolving sulphur in colourless $(NH_4)HS/NH_3$ solution. This solution is used to precipitate metal sulphides in qualitative analysis.

4. H_2S is a reducing agent and reduces Cl_2 to Cl^- , H_2SO_4 to SO_2 , $FeCl_3$ to $FeCl_2$ and itself gets oxidized to S .



Concept Ladder



H_2S is more acidic than H_2O and PH_3 . Thermal stability of H_2O is much higher than that of H_2S .

Rack your Brain



Why H_2S has greater molecular mass than H_2O , former is gas while latter is present in liquid form at room temperature?

Previous Year's Questions



Acidity of diprotic acids in aqueous solutions increases in the order

[NEET-2014]


- (1) $H_2S < H_2Se < H_2Te$
- (2) $H_2Se < H_2S < H_2Te$
- (3) $H_2Te > H_2S > H_2Se$
- (4) $H_2Se < H_2Te < H_2S$

[b] Sulphur Dioxide (SO₂) :

SO₂ is a colourless poisonous gas with a choking smell. It is highly soluble in water and almost completely present as SO₂·6H₂O in water and only 1% as H₂SO₃. Liquid SO₂ has also been used as a non-aqueous solvent.

Preparation:

1. It is prepared by burning S or H₂S in air.




2. It is also prepared by roasting various metal sulphides in smelters.




Chemical Properties :

1. Most of SO₂ is oxidized to SO₃ by the contact process and used to manufacture H₂SO₄.




2. Bleaching action:



Bleaching by SO₂ is temporary as reduced colour is reoxidised by air to give black coloured matter.

Also, H₂SO₄ produced during bleaching may be harmful to fibres.



Cl₂ although has a permanent effect as it produces [O].

Concept Ladder



H₂S acts only as a reducing agent while SO₂ can act both as a reducing and oxidising agent.

Rack your Brain



Why bleaching by SO₂ is temporary?

Previous Year's Questions





Nitrogen dioxide and sulphur dioxide have some properties in common. Which property is shown by these compounds, but not by the other?



[NEET-2015]

- (1) Is soluble in water.
- (2) Is used as a food preservative.
- (3) Forms 'acid-rain'.
- (4) Is a reducing agent.

3. It shows reducing properties in presence of moisture.




Although with stronger reducing agents it acts as an oxidising agent.



Detection of SO_2 :

It may be detected in three ways in the laboratory:

1. By its choking smell
2. It turns filter paper moistened with $\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+$ green.



Structure:

$3d_{z^2}$ orbital of S cannot be used to form π bond as its symmetry is wrong. Instead of $3d_{z^2}$ orbital on S is used to make the second π bond with $2p_x$ orbital on the second O atom. Although, it is surprising that both π bonds have the same energy.

Concept Ladder



Burning of sulphur or sulphide ores in air generates SO_2 .

Rack your Brain




What happens when SO_2 gas is passed through an aqueous solution of a Fe(III) salt.

Q.9 How SO_2 is an air pollutant?

A.9 SO_2 acts as an air pollutant because of the following reasons: (i) SO_2 is strongly irritating to the respiratory tract. SO_2 at a concentration of 5 ppm causes throat and eye irritation (resulting in cough, tears and redness in eyes). It causes breathlessness and affects larynx, i.e., voice box.

[c] Sulphur Trioxide (SO₃) :


It is the anhydride of H₂SO₄ and is solid at room temperature. It has three forms:



In gaseous state, SO₃ has a planar triangular structure involving sp² hybrid sulphur atom. The lone pair of sulphur atom forms coordinate bond (σ). Therefore, bond angle is exactly 120°. Solid SO₃ possesses either cyclic trimer (α – form) or infinite helical chains (β – form).


Preparation:

1. It is manufactured on a huge scale by the Contact Process. Practically all of it is converted to H₂SO₄ in the same process :




2. $\text{Fe}_2(\text{SO}_4)_3 \xrightarrow{\Delta} \text{Fe}_2\text{O}_3 + 3\text{SO}_3$


Chemical Properties:



SO₃ is used to prepare oleum (mainly pyrosulphuric acid H₂S₂O₇)



3. SO₃ is used to make sulfamic acid NH₂SO₃H (the only strong acid that can exist as a solid at room temperature).



Concept Ladder



- All the S—O bonds in SO₃ have equal length due to resonance.
- SO₃ exists in 3 allotropic forms α-SO₃, β-SO₃, γ-SO₃.

Rack your Brain



Why SO₃ is not absorbed directly in water to form H₂SO₄ in contact process?

Concept Ladder




SO₂ may be regarded as an anhydride of sulphurous acid (H₂SO₃) while SO₃ is regarded as an anhydride of sulphuric acid (H₂SO₄).

[d] Oxo Acids of Sulphur:

Some oxo acids of Sulphur ending in -ous have S in the oxidation state (+IV) and form salts ending in -ite. Some acid ending in -ic have S in the oxidation state (+VI) and form salts ending in -ate. Four main series exist which are:


(I) Sulphurous Acid Series:



The most important acid (according to the use) in this series is sulphurous acid (H_2SO_3):

Preparation:

It is only known in solution form (like carbonic acid) and as discussed SO_2 solution in water gives 1% H_2SO_3 .




Chemical Properties of SO_3^{2-} ion.


1. H_2SO_3 being dibasic forms salt of bisulphites (HSO_3^-) and sulphites (SO_3^{2-}). HSO_3^- however undergoes internal dehydration to form $\text{S}_2\text{O}_5^{2-}$ ion.



2. S(+IV) in SO_3^{2-} and HSO_3^- is a moderately strong reducing agent and forms sulphates on oxidation.



while with sulphur, sulphites form thiosulphates:



Rack your Brain



Define number of oxo acids formed by sulphure?

Concept Ladder



Sulphurous acid is an intermediate species in the formation of acid rain from sulphur dioxide.

Previous Year's Questions




Which of the following oxoacid of sulphur has —O—O— linkage

[NEET-2020]

- (1) H_2SO_3 , sulphurous acid
- (2) H_2SO_4 , sulphuric acid
- (3) $\text{H}_2\text{S}_2\text{O}_8$, peroxodisulphuric acid
- (4) $\text{H}_2\text{S}_2\text{O}_7$, pyrosulphuric acid




Sulphites and hydrogen sulphites liberates SO_2 on treatment with dilute acids:



3. SO_3^{2-} acts as an oxidising agent with a strong RA like H_2S .



(II) Sulphuric Acid Series:




(i) Sulphuric acid [Oil of Vitriol] H_2SO_4 :

It is the most important acid used in the chemical industry. It is a colourless syrupy liquid with B.P. 338°C and freezes into crystals at 10.5°C . It is strongly hydrogen bonded.


Preparation

1. In ancient days it was prepared from $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (Green vitriol).



2. Lead Chamber Process

Was used to manufacture H_2SO_4 industrially but the process is now obsolete as it only produced 78% H_2SO_4 . NO_2 is used as a homogeneous catalyst to oxidise SO_2 in the presence of water. NO produced combines with air to produce NO_2 which is reused. Thus, NO and NO_2 act as oxygen carriers



Concept Ladder



H_2SO_4 is considered as the **King of Chemicals**. In ancient days, it was called oil of vitriol (Green Vitriol).

Rack your Brain



Write down the main process for manufacturing of sulphuric acid.

Previous Year's Questions



Oleum is


[AIPMT]

- (1) castor oil
- (2) oil of vitriol
- (3) fuming H_2SO_4
- (4) none of these




3. Contact Process:

This process is most important commercial process for the manufacture of H_2SO_4 . It is used widely.



Flow diagram for the manufacture of sulphuric acid

- SO_2 is first collected from burning of sulphur or sulphide ores.
- SO_2 is then oxidized.



The forward rxn is exothermic and is favoured by high pressure, low temperature and excess of oxygen. A catalyst (activated V_2O_5) is used to obtain a reasonable conversion.

(iii) SO_3 is then passed into 98% H_2SO_4 forming pyrosulphuric acid ($\text{H}_2\text{S}_2\text{O}_7$) called oleum. Dilution of oleum with water gives concentrated H_2SO_4 (98%).

Previous Year's Questions




Identify the correct formula of oleum from the following :

[NEET-2019]

- (1) $\text{H}_2\text{S}_2\text{O}_7$
- (2) H_2SO_3
- (3) H_2SO_4
- (4) $\text{H}_2\text{S}_2\text{O}_8$

Chemical Properties :

1. H_2SO_4 due to its low volatility is used to manufacture more volatile acids.





2. H_2SO_4 is a powerful dehydrating agent especially for gases.

It dehydrates HNO_3 forming NO_2^+





It removes elements of H_2O from organic compounds.



H_2SO_4 mixes with H_2O to evolve large amount of heat. If H_2O is poured into concentrated acid, the heat evolved causes violent splashing. Thus, to dilute strong acids like H_2SO_4 , acid is poured into water while stirring (not water into acid).

3. H_2SO_4 is a quite strong oxidising agent.



4. Pure H_2SO_4 is a non-aqueous solvent and a sulphonating agent.



Q.10 In solution of H_2SO_4 when SO_3 is bubbled, a compound 'A' is formed, which further reacts with H_2O to give H_2SO_4 . Explain this process with equation?

A.10 H_2SO_4 absorbs SO_3 forming oleum (A) which reacts with H_2O to form H_2SO_4 .



Concept Ladder



- H_2SO_4 has high boiling point due to H-bonding. It acts as a strong dibasic acid.
- Metals those are more electropositive than hydrogen react with dil. H_2SO_4 to evolve H_2 gas but less electropositive metals on heating with conc. H_2SO_4 evolve SO_2 .

Rack your Brain



Why KMnO_4 should not be dissolved in conc. H_2SO_4 ?

Uses:

1. It is used in making superphosphate fertilizer.
2. It is used in removal of oxides from the surfaces of metals like Fe, Cu before electroplating or galvanizing.
3. It is used as a dehydrating agent, oxidising agent and as strong acid.


(ii) Thiosulphuric acid (H₂S₂O₃):

It is unstable and decomposes in H₂O into a mixture of S, H₂S, SO₂ and H₂SO₄ while the salts (thiosulphates) are stable. Large colourless hexagonal crystals are formed which are highly soluble in water.


sodium thiosulphate (Na₂S₂O₃·5H₂O) is the most important thiosulphate also called hypo.

Preparation:**Chemical Properties:**

1. Hypo solutions are used for iodine titrations in volumetric analysis.



In iodometric estimation oxidising agents like CuSO₄, K₂Cr₂O₇ are treated with KI which liberates I₂ which is determined by hypo as above.



2. Na₂S₂O₃ is used to destroy excess Cl₂ on fabrics after they have been bleached and called as antichlor.



It is also used to remove the taste from heavily chlorinated drinking water.

3. Hypo is used in photography for 'fixing' prints

Concept Ladder

H₂S₂O₃ is a dibasic acid. It does not exist in free state, salts are quite stable and are reducing in nature.

Rack your Brain

How many P—O—P bonds are there in (i) H₄P₂O₇, (ii) P₄O₁₀?


Previous Year's Questions

Which is the strongest acid in the following?


[NEET-2013]

- | | |
|------------------------------------|------------------------------------|
| (1) HClO ₄ | (2) H ₂ SO ₃ |
| (3) H ₂ SO ₄ | (4) HClO ₃ |


and films. It dissolves unreacted AgNO_3 , AgBr salts.



(iii) Thionic Acid Series:



Dithionic acid




Polythionic acid




(iv) Peroxo Acid Series

They contain the peroxo linkage $-\text{O}-\text{O}-$.



Peroxo monosulphuric acid
(Caro's Acid) H_2SO_5



Peroxo disulphuric acid $\text{H}_2\text{S}_2\text{O}_8$
(Marshall's Acid)

Concept Ladder



The chemical name for the following acids are: (i) H_2SO_5 Peroxymonosulphuric acid or Caro's acid, (ii) $\text{H}_2\text{S}_2\text{O}_8$ Peroxydisulphuric acid or Marshall's acid, (iii) $\text{H}_2\text{S}_2\text{O}_6$ Dithionic acid

Q.11 Why is sulphurous acid acts as reducing agent?

A.11 As Sulphur atom contains a lone pair of electrons, sulphurous acid can be easily oxidised to sulphuric acid. Therefore, it acts as a reducing agent.



Q.12 Write one chemical reaction to show that conc. H_2SO_4 can act as an oxidising agent.

A.12 Concentrated sulphuric acid oxidises carbon to CO_2 .





Group 17 (Halogen Family)

Introduction :

The elements Fluorine, Chlorine, Bromine, Iodine and Astatine constitute group 17 of the periodic table.

Electronic configuration :

Element	symbol	Atomic Number	Electronic Configuration	Brief representation of electronic configuration
Fluorine	F	9	$1s^2 2s^2 2p^5$	$[\text{He}] 2s^2 2p^5$
Chlorine	Cl	17	$1s^2 2s^2 2p^6 3s^2 3p^5$	$[\text{Ne}] 3s^2 3p^5$
Bromine	Br	35	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^5$	$[\text{Ar}] 3d^{10} 4s^2 4p^5$
Iodine	I	53	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^5$	$[\text{Kr}] 4d^{10} 5s^2 5p^5$
Astatine	At	85	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^5$	$[\text{Xe}] 4f^{14} 5d^{10} 6s^2 6p^5$

Occurrence :

1. Fluorine is less abundant in earth's crust than chlorine.
2. Fluorine is available mainly as insoluble fluorides.
3. Fluorine occurs to the extent of 0.07% of the earth's crust.

The important minerals of fluorine :

1. Fluorspar or fluorite : CaF_2
2. Cryolite : Na_3AlF_6
3. Fluorapatite: $3\text{Ca}_3(\text{PO}_4)_2 \cdot \text{CaF}_2$

Physical properties :

1. **Electronic configuration** : Their valence shell electronic configuration is $ns^2 np^5$ or $ns^2 np_x^2 np_y^2 np_z^1$ where $n = 2$ to 6 .
2. **Atomic and ionic radii** : The atomic radii increase from fluorine to iodine down the group due to increase in number of shells.

Concept Ladder



- Due to its very small size fluorine does not act as central atom in formation of interhalogen compounds.
- It is the most electronegative element in the whole periodic table.



3. **Ionization enthalpy** : Ionization energy of these elements are higher than those of the corresponding elements of group 16 due to increase nuclear charge.
4. **Electronegativity** : With increase in atomic number down the group, the electronegativity decreases. The decreasing order of electronegativity is $F > Cl > Br > I$.
5. **Electron gain enthalpy** : The decreasing order of negative electron gain enthalpy is $Cl > F > Br > I$.
6. **Oxidation state** : All the halogens show an oxidation state of -1 . Except fluorine, all halogens show positive oxidation state also.
7. **Metallic character** : All the halogens are non-metallic in nature due to their high ionization enthalpies and high electronegativities. The non-metallic character gradually decreases down the group.
8. **Bond dissociation energy** : Bond dissociation energy of fluorine is lower than those of chlorine ($Cl - Cl$) and bromine ($Br - Br$) because of inter electronic repulsions present in the small atom of fluorine. Hence, bond energy decreases in the order : $Cl_2 > Br_2 > F_2 > I_2$
9. **Bond length** : As the size of the halogen atom increases the bond length of $X - X$ bond in X_2 molecule increases from F_2 to I_2 .
10. **Melting point and boiling point** : It increases from F to I due to an increase in the VWF of attraction which increases down in the group.
11. **Density** : The density of halogen increases on moving down the group.
12. **Colour** : All halogens are coloured. The colour

Concept Ladder



Fluorine is better oxidising agent than chlorine, although it has less electron affinity. Chlorine has highest electron affinity in element in whole periodic table.

Rack your Brain



Why electron gain enthalpy of fluorine is less negative than that of chlorine?

Previous Year's Questions



Which one of the following orders is correct for the bond dissociation enthalpy of halogen molecules?

[NEET-2016]

- (1) $Br_2 > I_2 > F_2 > Cl_2$
- (2) $F_2 > Cl_2 > Br_2 > I_2$
- (3) $I_2 > Br_2 > Cl_2 > F_2$
- (4) $Cl_2 > Br_2 > F_2 > I_2$

darkness depends on the rise of atomic number from fluorine to iodine.

F : Light yellow; Cl : Yellow green;

Br : Raddish brown; I : Deep violet.

The colour is due to the adsorption of energy from visible light by their molecules to excite outer electrons for higher energy levels.

Chemical Properties

1. Reactivity :

All the halogens are chemically very reactive elements. This is due to their low dissociation energy. Fluorine is the most reactive and iodine is the least reactive halogen.

2. Thermal stability :

Thermal stability of hydrides decreases from HF to HI. i.e. $HF > HCl > HBr > HI$.

3. Oxidising power :

The oxidising power decreases on moving down the group. i.e. $F_2 > Cl_2 > Br_2 > I_2$.

Conversely, I^- is the strongest reducing agent while F^- is the weakest reducing agent.

4. Hydrogen halides :

All the halogens combine directly with H_2 to form covalent molecular hydrides but their reactivity progressively decreases from fluorine to iodine.

5. Bond strength, bond length and thermal stability:

Since, size of halogen atom increases from F to I down the group, bond length of H-X bond increases down the group.



Bond energy has inverse relationship with bond length i.e. larger the bond length, lower the bond strength and vice versa. Thus, order of bond strength is $HF > HCl > HBr > HI$

Bond strength provides a direct idea of thermal stability. Higher the bond dissociation

Concept Ladder



Standard reduction potential of halogens are positive and decreases down the group. Thus, halogens acts as strong oxidising agents and their oxidising power decreases from fluorine to iodine.

Previous Year's Questions



The variation of the boiling points of the hydrogen halides is in the order $HF > HI > HBr > HCl$.

What explains the higher boiling point of hydrogne fluoride?

[NEET-2015]

- (1) There is strong hydrogen bonding between HF molecules.
- (2) The bond energy of HF molecules is greater than in other hydrogen halides.
- (3) The effect of nuclear shielding is much reduced in fluorine which polarises the HF molecule.
- (4) The electronegativity of fluorine is much higher than for other elements in the group.

energy greater will be thermal stability. Thus, thermal stability follows the order.



5. **Reducing character :**


The reducing character of hydrogen halides increases down the group as, $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$



A less thermally stable compound has more tendency to release hydrogen easily and show greater reducing property. Since thermal stability of hydrogen halides decreases from HF to HI, their reducing power increases from HF to HI.

6. **Acidic strength :**

In gaseous state, hydrogen halides are covalent. But in aqueous solutions, they ionize and behave as acids. The acidic strength of these acids increases down the group and hence follows the order.



Reaction with H_2O :



Rack your Brain



Explain why halogens are strong oxidising agents.

Previous Year's Questions



Among the following, the correct order of acidity is

[NEET-2016]

- (1) $\text{HClO}_2 < \text{HClO} < \text{HClO}_3 < \text{HClO}_4$
- (2) $\text{HClO}_4 < \text{HClO}_2 < \text{HClO} < \text{HClO}_3$
- (3) $\text{HClO}_3 < \text{HClO}_4 < \text{HClO}_2 < \text{HClO}$
- (4) $\text{HClO} < \text{HClO}_3 < \text{HClO}_2 < \text{HClO}_4$

Q.13 Sea is the greatest source of halogens. Comment.

A.13 Sea water contains Cl, Br and I of Na, K, Mg and Ca, but mainly NaCl (2.5% by mass). Dried up sea beds contain NaCl and carnallite, $\text{KCl} \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$. Certain seaweeds contain upto 0.5% of iodine as sodium iodide and chile saltpetre (NaNO_3) contains upto 0.2% of sodium iodate. Therefore, sea is the greatest source of halogens.

- $3\text{H}_2\text{O} + 3\text{F}_2 \longrightarrow 6\text{HF} + \text{O}_3$
- $\text{Cl}_2 + \text{H}_2\text{O} \longrightarrow \text{HCl} + \text{HClO}$
(Hypochlorous acid)
- $\text{Br}_2 + \text{H}_2\text{O} \longrightarrow \text{HBr} + \text{HBrO}$
(Hypobromous acid)
- $2\text{HClO} \longrightarrow 2\text{HCl} + \text{O}_2$
 $2\text{HBrO} \longrightarrow 2\text{HBr} + \text{O}_2$
- $\text{I}_2 + \text{KI} = \text{KI}_3 + \text{I}^- = \text{I}_3^-$

Reaction with hydrogen :

- $\text{H}_2 + \text{F}_2 \xrightarrow{200^\circ\text{C}} 2\text{HF}$ (very violent)
 $\text{H}_2 + \text{Cl}_2 \xrightarrow{\text{Sunlight}} 2\text{HCl}$
 $\text{H}_2 + \text{Br}_2 \xrightarrow[\text{Pt. catalyst}]{\text{Heat}} 2\text{HBr}$
 $\text{H}_2 + \text{I}_2 \xrightleftharpoons{\text{heat}} 2\text{HI}$ (Poor yield)
- Bleaching action of halogen : Cl_2 acts as a bleaching agent. Its bleaching action is permanent Cl_2 water can also acts as ink remover.
- Reaction with other halides :
 $2\text{KBr}(\text{aq}) + \text{Cl}_2(\text{g}) \longrightarrow 2\text{KCl}(\text{aq}) + \text{Br}_2(\text{g})$
 $2\text{KI}(\text{aq}) + \text{Cl}_2(\text{g}) \longrightarrow 2\text{KCl}(\text{aq}) + \text{I}_2(\text{g})$

Anomalous behaviour of fluorine :

Fluorine differs from other members of its own sub-group due to the following reasons.

- Small size of the atom in comparison to the sizes of Cl, Br and I atoms.
- Higher electronegativity.
- Absence of d-orbitals in its valence shell.
- Higher positive reduction potential.

The main points of difference are :

- Fluorine shows only a negative oxidation state

Concept Ladder



Among halogens, fluorine provides the largest variety of inter-halogen compounds.

Rack your Brain



Arrange hydrogen halides in the increasing order of their dipole moments.

Previous Year's Questions



Which of the following statements is not true for halogens?

[NEET-2018]

- All form monobasic oxyacids
- All are oxidizing agents.
- All but not fluorine, show positive oxidation states.
- Chlorine has the highest electron-gain enthalpy.



Reaction with			
F ₂	Cl ₂	Br ₂	I ₂
Preparation Of Gas			
<p>Light - yellow gas</p> <p>Most reactive element known strong O.A.</p> <p>Moisson was awarded Noble prize for his methods of preparing F₂ :</p> $\text{CaF}_2 + \text{conc. H}_2\text{SO}_4 \longrightarrow \text{HF}$ $\text{HF} + \text{KF} \longrightarrow \text{K}^+ [\text{HF}_2]^-$	<p>Yellow green gas</p> <p>toxic, has pungent suffocating odour</p> <p>Lab</p> $\text{MnO}_2 + 4\text{HCl} \longrightarrow \text{MnCl}_2 + 2\text{H}_2\text{O} + \text{Cl}_2$ <p>Deacon's Process :</p> <p>It is them dried with CaO, P₂O₅ or H₂SO₄.</p> $\text{HCl} + \text{O}_2 \xrightarrow{\text{CuCl}_2} \text{Cl}_2 + \text{H}_2\text{O}$ <p>Commercially it is also obtained by electrolysis of NaCl(aq) in the manufacture of NaOH or elect. of NaCl(l) in the manufacture of Na</p>	<p>Brown liquid</p> <p>Obtained from Sea water and brine lakes by passing Cl₂ as follows:</p> $\text{Cl}_2 + 2\text{Br}^- \longrightarrow \text{Cl}^- + \text{Br}_2$ $\text{Br}_2 + 3\text{Na}_2\text{CO}_3 \longrightarrow \text{NaBr} + \text{NaBrO}_3 + 3\text{CO}_2$ $\text{Br}_2 \xleftarrow{\text{CuCl}_2} \text{HBr} + \text{HBrO}_3$	<p>Violet solid</p> <p>Crystallizes as black flakes has slight lustre.</p> <p>Chile saltpetre is mainly NaNO₃, but contains traces of NaIO₃, NaNO₂, is crystallised out and left NaIO₃ is divided into two parts</p> $\text{NaIO}_3 \xrightarrow{\text{NaHSO}_3} \text{NaIO}_2/\text{H}^+ + \text{NaHSO}_4$ $\text{NaIO}_2/\text{H}^+ \xrightarrow{\text{I}^- + \text{SO}_4^{2-}} \text{I}_2(\text{solid})$ <p>Purification is done by sublimation.</p>
H ₂ O			
$\text{F}_2 + \text{H}_2\text{O} \longrightarrow \text{2H}^+ + 2\text{F}^- + \frac{1}{2} \text{O}_2$ <p>(OA)</p> <p>(spontaneous and strongly exothermic)</p>	$\text{Cl}_2 + \text{H}_2\text{O} \xrightleftharpoons[\text{K}^+]{\text{Cold OH}^-} \text{ClO}^- + \text{Cl}^-$ $\text{Cl}_2 + 3\text{H}_2\text{O} \xrightarrow{\text{hot OH}^-} \text{ClO}_3^- + \text{Cl}^-$	$\text{Br}_2 + \text{H}_2\text{O} \xrightarrow{\text{OH}^-} \text{Br}^- + \text{BrO}^-$	<p>Sparingly soluble in water but dissolves if I⁻ are present (as soluble I₃⁻ forms)</p> $\text{I}^- + \text{I}_2 \longrightarrow \text{I}_3^-$
NH ₃			
X ₂ reacts with NH ₃ to give NH ₃ X and NH ₄ X (X= F, Cl, Br)			
Others			
All halogens oxidise H ₂ S to S ; form SO ₂ X ₂ with SO ₂ ; trihalides with As, Sb, Bi; penta halides (except I) with As, Sb, Bi .			
Uses			
Used in making freon (CFC's)	Used as a bleaching agent $\text{Cl}_2 + \text{H}_2\text{O} \longrightarrow \text{HOCl}$ and make poisonous gases like SO ₂ Cl ₂	It is used to make flame retardant and in organic chemistry.	It is used to make tincture of iodine (I ₂ + KI) and in titrations.

of -1 due to maximum value of electronegativity. It never shows any positive oxidation state. The other halogen show negative as well as positive oxidation states, i.e. between -1 and $+7$.

2. Fluorine is most reactive as bond dissociation energy of $F - F$ bond is low. $X - X$ bond is stronger in Cl_2 and Br_2 .

3. Maximum covalency of fluorine is one as there is no d -orbital in its valence shell. Other members can have maximum covalency of 7 because of vacant d -orbitals.

4. Fluorine does not have vacant d -orbital in valence shell therefore it does not combine with F^- ion to form polyfluoride ions like Cl^- , Br^{2-} , I_6^- etc.

Compounds of Halogen :

Chlorine: (Cl_2)

Occurrence :

1. Chlorine is widely distributed in nature in the combined state in the form of chlorides of various metals.

2. Common Salt ($NaCl$) is the most important chloride which occurs in sea water, lakes and in rocks.

The other important chloride minerals are :

1. Sylvine (Potassium Chloride) KCl
2. Carnalite, $KCl, MgCl_2, 6H_2O$
3. Chlorapatite, $3Ca_3(PO_4)_2, CaCl_2$
4. Horn Silver, $AgCl$

Preparation :

1. By oxidation of conc. HCl

Concept Ladder



Fluorine and oxygen compounds are called oxygen fluorides, but not fluorine oxides.

Rack your Brain



Why I_2 is more soluble in KI than in water?


Previous Year's Questions




Which of the following elements is extracted commercially by the electrolysis of an aqueous solution of its compound?

[AIPMT]

- | | |
|----------|----------|
| (1) Cl | (2) Br |
| (3) Al | (4) Na |




2. Weldon's process : By heating pyrolusite with conc. HCl



3. Deacon Process :

HCl(g) is oxidised by atm. oxygen in presence of a catalyst CuCl_2 at 400°C (723 K).



Physical Properties :

It is yellowish green gas, poisonous in nature, soluble in water. Its aqueous solution is known as chlorine water which on careful cooling gives chlorine hydrate $\text{Cl}_2 \cdot 8\text{H}_2\text{O}$.

Chemical Properties :

1. Bleaching action and oxidising property.



The bleaching action of chlorine is permanent and is due to its oxidising nature.



4. Action of hydrogen :



5. Displacement reactions :



6. Action of NaOH :



Concept Ladder



Deacon's process is superior to Weldon's process as in Deacon's process, chlorine is obtained by the oxidation of HCl with air in presence of cupric chloride heated at 400°C .

Rack your Brain



What is the cause of bleaching action of chlorine in water?

Previous Year's Questions



The correct order of increasing bond angles in the following species is

[AIPMT]

- (1) $\text{Cl}_2\text{O} < \text{ClO}_2 < \text{ClO}_2^-$
- (2) $\text{ClO}_2 < \text{Cl}_2\text{O} < \text{ClO}_2^-$
- (3) $\text{Cl}_2\text{O} < \text{ClO}_2^- < \text{ClO}_2$
- (4) $\text{ClO}_2^- < \text{Cl}_2\text{O} < \text{ClO}_2$



7. Combination with non metals



8. Other important reactions

Reaction with ammonia



Reaction with hydrocarbons



Oxidizing nature of chlorine



Uses of Chlorine :

It is used as :

1. Bleaching agent
2. Disinfectant
3. In the manufacture of CHCl_3 , CCl_4 , DDT, BHC. (Insecticides) etc.
4. In the manufacture of synthetic plastic such as PVC.
5. Poisonous gas phosgene (COCl_2), tear gas (CCl_3NO_2), Mustard gas ($\text{ClC}_2\text{H}_4\text{SC}_2\text{H}_4\text{Cl}$).

Concept Ladder



Bleaching of flowers by Cl_2 is permanent, while that by SO_2 is temporary.

Rack your Brain



Arrange the following oxoacids of chlorine in increasing order of their acidic strength.

HOCl , HOClO , HOClO_2 , HOClO_3

Previous Year's Questions



Which of the following displaces Br_2 from an aqueous solution containing bromide ions?

[AIPMT]

- | | |
|-------------------|--------------------|
| (1) I_2 | (2) I_2^- |
| (3) Cl_2 | (4) Cl^- |

- Refrigerant such as freon (CCl_2F_2).
- Extraction of Metals like Au and Pt.

Hydrogen Chloride (Hydrochloric Acid) HCl :

Preparation :

- HCl is prepared by dissolving hydrogen chloride gas in water.
- Hydrogen chloride gas required in turn can be prepared by the following methods.



HCl gas can also be obtained by burning hydrogen in chlorine.

Laboratory Method :

- In the laboratory HCl is prepared by mixture of NaCl and conc. H_2SO_4



- $\text{HCl}(\text{g})$ is dried by passing through cone, H_2SO_4 collected by upward displacement of air.

Properties :

- HCl is a covalent compound but when dissolved in H_2O it ionizes to form hydrogen ions or chloride ions or hydronium ions.



- It reacts with metal to form their respective chlorides and $\text{H}_2(\text{g})$.



Reaction with NH_3 :



- It reacts with metal carbonates, bicarbonates, sulphides, sulphites, thiosulphates and nitrites

Concept Ladder



HCl also known as muriatic acid, is an aqueous solution of hydrogen chloride. It is a colourless solution with a distinctive pungent smell.

Rack your Brain



Explain why ClF_3 exists but FCl_3 does not?

Previous Year's Questions





Elements of which of the following groups will form anions most readily?

[AIPMT]

- Oxygen family
- Nitrogen family
- Halogens
- Alkali metals

etc to form their respective chlorides.



2. It is used for dissolving noble metals like Au, Pt etc.



Uses :


1. In the preparation of chlorides, chlorine, aqua-regia etc.
2. To manufacture glucose from corn starch.
3. In medicine and in galvanizing.
4. In the production of dyes, paints, photographic chemicals etc.

Interhalogen Compounds :

1. These compounds are regarded as halides of more electropositive (i.e. less electronegative) halogens.

Q.14 How can you prepare Cl_2 from HCl and HCl from Cl_2 . Write reactions only.

A.14 HCl can be oxidised to Cl_2 by a number of oxidising agents such as MnO_2 , KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$, etc.



Cl_2 can be reduced to HCl by its reaction with H_2 in presence of diffused sunlight.



Concept Ladder



As chlorine is second most electronegative element in the periodic table, it can form compounds as carbonates, bicarbonates, sulphides and nitrates.

Rack your Brain



Write the components used in the formation of aquaregia.



HOCI/HClO Hypochlorous acid	HClO ₂ Chlorous acid	HClO ₃ Chloric acid	HClO ₄ Perchloric acid
General characteristics			
Very weak acid Only known in aqueous solution. NaOCl is well known Unstable	weak acid (stronger than HClO) Only exists in solution.	Known as salts and in aqueous Not very stable and detonates if heated $2\text{HClO}_3 \rightarrow 2\text{ClO}_2 + \text{H}_2\text{O} + 1/2\text{O}_2$	One of the strongest acids known. Both salts and acids are well known, can in fact be isolated in anhydrous form ($\text{HClO}_4 \cdot 2\text{H}_2\text{O}$)
Preparation			
Shake halogen with freshly precipitated HgO $2\text{HgO} + \text{H}_2\text{O} + 2\text{Cl}_2 \rightarrow \text{HgO} \cdot \text{HgCl}_2 + 2\text{HOCl}$	$\text{Ba}(\text{ClO}_2)_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 \downarrow + \text{H}_2\text{ClO}_2$ BaSO ₄ is filtered off	$\text{Ba}(\text{ClO}_3)_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2\text{HClO}_3$ BaSO ₄ is filtered off	$\text{Ba}(\text{ClO}_4)_2 + \text{H}_2\text{SO}_4 \rightarrow \text{BaSO}_4 + 2\text{HClO}_4$
NaOCl is made industrially by electrolysis of cold brine (NaOH solution) or by: $1/2\text{X}_2 + \text{OH}^- \xrightarrow[\text{<RT}]{\text{Cold}} \text{XO}^- + \text{HX}$	Chlorites are made from ClO ₂ and NaOH ClO ₂ and Na ₂ O ₂ (See Chlorine Dioxide)	Chlorates are formed by: $\text{X}_2 + \text{OH}^- \xrightarrow{\text{Hot}} \text{XO}_2^- + \text{X}^-$	$\text{NaClO}_2 + \text{H}_2\text{O} \xrightarrow{\text{Electrolysis}} \text{NaClO}_4 + \text{H}_2$
Reactions			
$3\text{XO}^- \xrightarrow{\text{hot}} 2\text{XO}_2^- + \text{X}^-$ +1 +V +1	$\text{ClO}_2 \xrightarrow{\text{OH}^-} \text{stable}$	$\text{ClO}_2 \xrightarrow{400^\circ\text{C}} \text{stable}$	$2\text{HClO}_4 \xrightarrow[-\text{H}_2\text{O}]{\text{P}_2\text{O}_5} \text{Cl}_2\text{O}_7$
$2\text{HClO} \xrightarrow{-\text{H}_2\text{O}} \text{Cl}_2\text{O}$	$2\text{ClO}_2 \xrightarrow[\Delta]{2\text{OH}^-} \text{ClO}_2^- + \text{ClO}_3^- + \text{H}_2\text{O}$ makes solution yellow	$\text{ClO}_2 \xrightarrow[\text{low temp.}]{\Delta} \text{ClO}_2^- + \text{Cl}^-$	$2\text{HClO}_3 \xrightarrow{-\text{H}_2\text{O}} \text{Cl}_2\text{O}_7$
Uses			
Used in bleaching and as a disinfectant (especially NaOCl)	Chlorites are also used as bleaches	Chlorates are used to make fire works, matches, NaClO ₃ is a powerful weedkiller: Solid NaClO ₃ has also been used to make bombs.	Half of perchlorate is used as NH ₄ ClO ₄ in shuttle launch. KClO ₄ is used in fireworks and flares.

2. Types of interhalogen compound :

AB type : ClF, BrF, BrCl, ICl, IBr

AB₃ type : ClF₃, BrF₃, ICl₃

AB₅ type : BrF₅, IF₅

AB₇ type : IF₇

Preparation :

- $\text{Cl}_2 + \text{F}_2 \xrightarrow{437\text{K}} 2 \text{ClF}$
- $\text{Cl}_2 + 3\text{F}_2 \xrightarrow{573\text{K}} 2 \text{ClF}_3$
- $\text{I}_2 + 3\text{Cl}_2 \longrightarrow 2 \text{ICl}_3$
- $\text{Br}_2 + 3\text{F}_2 \longrightarrow 2 \text{BrF}_3$
- $\text{I}_2 + \text{Cl}_2 \longrightarrow 2 \text{ICl}$
- $\text{Br}_2 + 5\text{SF}_2 \longrightarrow 2 \text{BrF}_5 + 5\text{S}$


Structure :

1. AB Type :

The central halogen, i.e., the bigger halogen atom (A) is sp³ hybridised. The molecules are linear due to the presence of three lone pairs on A atom.

2. AB₃ Type :

The central halogen atom is sp³d hybridised. The molecules have a bent T-shaped geometry instead of trigonal bipyramidal due to the presence of two lone pairs.



Structure of BrF₃

3. AB₅ Type :

The central halogen atom is sp³d² hybridised leading to a square pyramidal geometry instead of octahedral geometry as there present 1 lone pair on the halogen atom A.

Concept Ladder



In general interhalogen compounds are more reactive than pure halogens except fluorine because A—B bond is weaker than B—B bond.

Rack your Brain




Why interhalogen compounds are strong oxidizing agents.

Concept Ladder




Four types of interhalogen compounds are :

- XX' : ClF, BrF, BrCl etc.
- XX'₃ : ClF₃, BrF₃, IF₃ etc.
- XX'₅ : ClF₅, BrF₅, IF₅ etc.
- XX'₇ : IF₇



4. AB₅ Type :

The central halogen atom (A) is sp^3d^2 hybridised and the molecules shows the expected pentagonal bipyramidal shape since there are no lone pairs.



Characteristics of Interhalogen Compound :

1. These may be gases (ClF , BrF , ClF_3 , IF_7), liquids (BrF_3 , BrF_5 , IF_5) or solids (ICl , IBr , IF_3 , ICl_3).
2. The interhalogen molecules are diamagnetic in nature.
3. With increase in ΔEN between A and B, thermal stability of AB type compounds increases. The more polar is the A – B bond and hence greater is the thermal stability.



4. A_2 or B_2 molecules are less reactions than AB type compounds. Since AB bond is weaker than A–A & B–B bonds. These order of reactivity of some interhalogen compounds has been found as : $ClF_3 > BrF_5 > IF_7 > BrF_3 > IF_5 > BrF$

6. Hydrolysis gives halogen acid and oxyhalogen acid. The oxyhalogen acid is of larger halogen atom.



Concept Ladder



The interhalogen compounds of iodine and fluorine are thermally more stable while the compounds of iodine and chlorine or iodine and bromine are less stable.

Rack your Brain



Write the general formula used for formation of interhalogen compounds.

Concept Ladder



AB type interhalogen compounds are ClF , BrF , $BrCl$, ICl , IBr .



7. The interhalogens of AB type add at olefinic double bond sites.



Uses Of Interhalogen Compounds :

1. ClF_3 and BrF_3 are also used as fluorinating agents.
2. Interhalogen compound are used for the preparation of polyhalides.
3. ClF_3 and BrF_3 are used for the production of UF_6 in the enrichment of ^{235}U .



4. Interhalogen compounds can be used as non-aqueous solvents. BrF_3 is also used as Lewis acid.
5. Interhalogen compounds of the type XX' (particularly ICl) are used as halogenating agent. ICl is used in the estimation of iodine number of fats and oils.

Concept Ladder



Interhalogen compounds are gases or liquids, volatile and create fume in air. These can be hydrolysed and act as oxidising agents.

Rack your Brain



What are polyhalides? Write some examples of polyhalides in which 2 or 3 different halogens are present.











Q.15 Why interhalogens are more reactive than halogens?

A.15 This is because the bond in the interhalogen ($\text{X}-\text{X}'$) is weaker than $\text{X}-\text{X}$ and $\text{X}'-\text{X}'$ bond in the halogens. This is due to less effective overlapping between orbitals of dissimilar atoms than those between similar atoms. However, due to low bond dissociation enthalpy of $\text{F}-\text{F}$ bond, F_2 is more reactive than its interhalogen compounds.

Oxyacids or Oxoacids of Halogens :

Halogen	Hypohalous acids (x = +1)	Halous acid(x = +3)	Halic acid (x = +5)	Perhalic acid (x = +7)
Cl	HClO	HClO ₂	HClO ₃	HClO ₄
Br	HBrO	—	HBrO ₃	—
I	HIO	—	HIO ₃	HIO ₄

Structure : Central halogen atom of all oxyacids of halogen is sp³ hybridised.



Oxidising power of these oxyacids decreases as the oxidation number of halogen increases.




Thermal stability increases with increase in oxidation state of halogen.



Hydra – Acids

I Hydrofluoric acid [H_2F_2 or HF] :

1. HF is a colourless, corrosive liquid with pungent smell. It attacks glass and is thus stored in wax bottles or gutta percha bottles.



(This is also used for etching glass.)

HF is kept in laboratory in KF giving K^+ $[\text{F}-\text{H}-\text{F}]^-$ ions.

2. H_2 and F_2 combine with each other very violently even in dark to form HF. Thus special methods are employed in its preparation.

Industrially it is made by heating CaF_2 with strong H_2SO_4 . The reaction is endothermic, hence the need for heating as SiO_2 impurities are removed from CaF_2 otherwise they consume much of HF produced.



3. Two thirds of HF produced are used to make chlorofluoro carbons (Freons) used as refrigerating fluids.



Concept Ladder



Unlike other halogen acids, hydro fluoric acids attacks silica and glass. With silica it forms silicon tetrafluoride and hydro thio salicylic acid.

Rack your Brain



Why on heating H_2F_2 with a mixture of MnO_2 and H_2SO_4 no gas is involved?

Previous Year's Questions




Which of the statements given below is incorrect?

[NEET-2015]


- (1) O_3 molecule is bent.
- (2) ONF is isoelectronic with O_2N^- .
- (3) OF_2 is an oxide of fluorine.
- (4) Cl_2O_7 is an anhydride of perchloric acid.

4. In liquid form HF is used as a non - aqueous solvent.



II. Hydrochloric acid [HCl]


1. At one time HCl was made exclusively by 'salt cake' method.




2. Highly pure HCl is made by bringing H_2 and Cl_2 in a special combustion chamber. (Direct combination is explosive)



3. In laboratory :



4. Cl cannot be dried over P_2O_5 or quick lime because it reacts with them:



Hence it is dried by passing through conc. H_2SO_4

5. HCl(g) is colourless, weak reducing agent (is oxidized by strong oxidising agents like MnO_2 , KMnO_4 , $\text{K}_2\text{Cr}_2\text{O}_7$) and highly soluble in water.

III HBr and HI

1. They are prepared by reaction of phosphoric acid on metal bromides or iodides (similar to salt cake process)



Concept Ladder



HBr and HI cannot be prepared by heating bromides and iodides with concentrated H_2SO_4 while HCl can.

Rack your Brain



How HCl is considered to be strong acid.

Previous Year's Questions





Bleaching powder reacts with a few drops of conc. HCl to give


[AIPMT]

- (1) chlorine
- (2) hypochlorous acid
- (3) calcium oxide
- (4) oxygen

2. The usual laboratory process is :




3. HBr is not very stable and acts as a strong reducing agent while HI is least stable and thus a more powerful reducing agent.



Oxides

The bonds in oxides are largely covalent as there is only a small difference in electronegativity between the halogens and oxygen. Most halogen oxides are unstable and tend to explode when subjected to shock. Stability of oxides increases down the group and higher oxidation states are more stable than lower oxidation states.

I Dichloro oxide [Cl_2O]



Cl_2O is a yellow – brown gas which condenses to an orange liquid (bp = 2°C). It is the anhydride of hypochlorous acid (HClO).

1. Cl_2O is prepared by passing dry chlorine over freshly precipitated mercuric oxide.



2. It is soluble in water



Concept Ladder



HF is the weakest acid among hydro halo acids.

HF < HCl < HBr < HI

Rack your Brain



With what neutral molecule ClO^- is isoelectronic?

Previous Year's Questions




The correct order of increasing bond angles in the following species is :

[AIPMT]

- (1) $\text{Cl}_2\text{O} < \text{ClO}_2 < \text{ClO}_2^-$
- (2) $\text{ClO}_2 < \text{Cl}_2\text{O} < \text{ClO}_2^-$
- (3) $\text{Cl}_2\text{O} < \text{ClO}_2^- < \text{ClO}_2$
- (4) $\text{ClO}_2^- < \text{Cl}_2\text{O} < \text{ClO}_2$


3. It explodes in presence of NH_3 or a reducing agent or on heating.



4. It is a strong oxidising agent.



II Chlorine Dioxide [ClO_2]



It is a yellow gas, condenses to deep red liquid, explodes above -40°C and in presence of reducing agents. It is paramagnetic being an odd electron molecule but does not dimerise as the odd electron is delocalised.

1. It explodes when mixed with reducing agents. The safest laboratory preparation is from sodium chlorate and oxalic acid as this automatically dilutes the gas with CO_2 .



2. $2\text{ClO}_2 + \text{H}_2\text{O} \longrightarrow \underbrace{\text{HClO}_3 + \text{HClO}_2}_{\text{dark green solution}} + \text{heat}$

3. It is used to manufacture NaClO_2 which is used for bleaching paper textile.



4. ClO_2 is a powerful oxidising agent and a bleaching agent and is used to purify water.

Concept Ladder



ClO_2 is a powerful oxidising and bleaching agent. It has an angular structure with $\text{O}-\text{Cl}-\text{O}$ bond angle of 117.6° . The molecule is supposed to contain a 3 electrons bond.


Rack your Brain




Write equation when ClO_2 reacts with alkalis.

Q.16 What happens when Cl_2 is passed through a hot concentrated solution of a base like $\text{Ba}(\text{OH})_2$?

A.16 On treatment with $\text{Ba}(\text{OH})_2$, Cl_2 undergoes disproportionation to form barium chloride, $\text{Ba}(\text{ClO}_3)_2$.




III Dichlorine Hexoxide [Cl₂O₆]




The exact structure is unknown

It is a dark red liquid which freeze at -180°C to a yellow solid. It is diamagnetic in nature due to absence of unpaired electrons and exists as a dimer. The structure is guessed to be as shown but the exact structure is unknown.

1. It is obtained from ClO_2 and O_3 .



2. Hydrolysis with water or alkali gives chlorate and perchlorate




3. It is a strong oxidising agent which explodes on contact with grease.

4. Reaction with anhydrous HF is reversible:



IV Dichlorine Heptoxide [Cl₂O₇]

Cl_2O_7 is a colourless oily liquid which is moderately stable and less reactive than others. It is the only exothermic oxide of Cl and the anhydride of perchloric acid HClO_4 . It is anhydride of perchloric acid and made by dehydrating perchloric acid with P_2O_5 or H_3PO_4 .



Concept Ladder



In vapour state, Cl_2O_6 exists as ClO_3 molecule which is para magnetic while in liquid state it is in dimeric form having even no. of electron hence diamagnetic in nature.

Previous Year's Questions



Which one of the following oxides is expected to exhibit paramagnetic behaviour?

[AIPMT]


- (1) CO_2 (2) SiO_2
(3) SO_2 (4) ClO_2

Bleaching Powder:

Bleaching powder is actually written as $\text{Ca(OCl)}_2 \cdot \text{Ca(OH)}_2 \cdot \text{CaCl}_2 \cdot 2\text{H}_2\text{O}$. It is a pale-yellow powder having a strong smell of chlorine and soluble in water. However, a clear solution is never formed due to the presence of impurities.



Preparation:

It is made by passing Cl_2 into slaked lime.



Chemical Properties:

1. Bleaching action



2. Oxidising action



3. It converts acetone into chloroform.

Concept Ladder



Bleaching powder is also termed as calcium chloro hypochlorite, because it is considered as a mixed salt of hydrochloric acid and hypochlorous acid.

Rack your Brain



Name the processes used for manufacturing of bleaching powder?

Q.17 Halogens have maximum negative electron gain enthalpy. Explain why?

A.17 Halogen have high effective nuclear charge due to its smallest size in their respective periods. Therefore, they readily accept one electron to acquire the stable electronic configuration of the nearest noble gas. In other words, a halogen atom can accepts an electron therefore larger amount of energy is released, thus halogens have maximum negative electron gain enthalpies.



Group 18 (Noble gases)

Introduction :

1. The elements Helium, Neon, Argon, Krypton, Xenon and Radon constitute group 18 (or zero) of the periodic table.
2. All the noble gases, except He have eight electrons (closed shell) with electronic configuration ns^2np^6 where $n = 2$ to 6. Helium, however has only two electrons and hence its electronic configuration is $1s^2$.

Electronic Configuration :

Element	Symbol	Atomic Number	Electronic Configuration
Helium	He	2	$1s^2$
Neon	Ne	10	$1s^2 2s^2 2p^6$ OR [He] $2s^2 2p^6$
Argon	Ar	18	$1s^2 2s^2 2p^6 3s^2 3p^6$ OR [Ne] $3s^2 3p^6$
Krypton	Kr	36	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6$ OR [Ar] $3d^{10} 4s^2 4p^6$
Xenon	Xe	54	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6$ OR [Kr] $4d^{10} 5s^2 5p^6$
Radon	Rn	86	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 6s^2 6p^6$ OR [Xe] $4f^{14} 5d^{10} 6s^2 6p^6$

Occurrence :

1. Except radon (radioactive) all noble gases are present in the atmosphere throughout the universe in the free state to the extent of 1% by volume. Helium is also found as natural gas **Ramsay** discovered. He in cleveite (a uranium ore) i.e. on the earth.
2. The gas is obtained when these minerals are either heated at 1273 K in vacuum or heated with H_2SO_4 . Helium and argon are also found in the dissolved gases of some mineral.

Concept Ladder



The zero group occupies the intermediate position between the elements of VII A and I A groups.

Rack your Brain



Why noble gases are monoatomic in nature?

3. Argon is the most abundant noble gas in the atmosphere (0.934% by volume).

The relative abundances of other gases in the atmosphere are :

$$\text{He} = 5.24 \times 10^{-4}, \text{Ne} = 1.82 \times 10^{-3}$$

$$\text{Ar} = 93.4 \times 10^{-2}, \text{Kr} = 1.45 \times 10^{-3}$$

$$\text{and Xe} = 8.70 \times 10^{-6} \% \text{ by volume}$$

Physical Properties of Noble Gases

1. Physical state :

All the elements of group 18 are colourless, odourless and tasteless.

2. Atomic radii :

The atomic radii of noble gases increases on moving down-the group and their atomic radii correspond to the vander waal's radii.

$$\text{Order : He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe} < \text{Rn}$$

3. Melting point and boiling point :

M.P. and B.P. increases from He to Rn, because of increase in magnitude of vander waal's forces.

$$\text{Order : He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe} < \text{Rn}$$

4. Ionisation energy and electron gain enthalpy :

Noble gases have stable $ns^2 np^6$ fully filled electronic configuration so, these have no tendency to add or lose electron. Therefore ionisation energy of noble gas is very high. On the other hand their electron gain enthalpy is zero.

$$\text{Order : He} > \text{Ne} > \text{Ar} > \text{Kr} > \text{Xe} > \text{Rn}$$

5. Solubility in water :

They are slightly soluble in water their solubility generally increases with the increase in atomic number down the group.

$$\text{Order : He} < \text{Ne} < \text{Ar} < \text{Kr} < \text{Xe} < \text{Rn}$$

Concept Ladder



Noble gases are also termed as inert gases or the rare gases of the atmosphere.

Rack your Brain



Explain why electron gain enthalpy of noble gases is zero?

Concept Ladder



Noble gases have lowest boiling point in comparison to the elements of their respective periods.

6. Liquification of gases :

It is difficult to liquify inert gases as their atoms are held by weak VWF.

7. Ease of liquification increases from He to Rn.

8. He has the lowest B.P. (4.18 K) of any known substances. The ease of liquification increases from He to Rn due to increase in inter-molecular forces.

9. The zero group of the periodic table constitutes Helium (He), Neon (Ne), Argon (Ar), Krypton (Kr), Xenon (Xe) and Radon (Rn).

10. Noble gases do not have chemical reactivity and at ordinary temperature they are present in gaseous form and therefore, these are called **Inert gases**.

Order : He < Ne < Ar < Kr < Xe < Rn

11. Ability to diffuse :

All the noble gas are able to diffuse through glass, rubber, plastic, material and some metals.

Order : He > Ne > Ar > Kr > Xe > Rn

Uses Of Noble Gases :

Helium :

1. Helium is a non-inflammable and light gas therefore, it is used in filling balloons for meteorological observations.

2. It is also used in gas cooled nuclear reactors.

3. Liquid helium (boiling point 4.2 K) is used as Cryogenic agent for carrying out various experiments at low temperatures.

Neon :

1. Ne is used in fluorescent bulbs and discharge tubes for advertisement display purposes.

Concept Ladder



Helium forms no real chemical compounds. Among noble gases only Xenon forms chemical compounds.

Rack your Brain



Noble gases can form compounds only with two elements. Name them?

Concept Ladder



Helium — oxygen mixture is used for artificial respiration in deep seas diving and in treatment of asthma.

2. Neon is also used in televisions set, spark plug, warning signals etc.
3. Glow of different coloured 'neon signs' can be produced by mixing neon with other gases.

Argon :

1. Ar is mainly used to provide an inert atmosphere high temp. metallurgical processes (alloys or arc welding of metals) and for filling electric bulbs. It is also used in the laboratory to handle substances that are air sensitive.
2. This gas is used in gas chromatography.
3. It is mixed with neon to obtain different colours in neon sign lamps.

Krypton And Xenon :

1. Kr and Xe are used in light bulbs designed for special purpose.
2. These are also used for high speed photography.
3. Krypton-85 is used to measure thickness of sheets of metals and plastics.

Radon :

1. Radon being radioactive is used in nuclear processes.
2. It is also used in treatment of cancer and other tumors.
3. It is used for detecting dislocations and defects in metals and other solids.

Concept Ladder

The two methods used for preparation of argon gas is Ramsay method and Rayleigh method.

Previous Year's Questions

Identify the incorrect statement, regarding the molecule XeO_4 :

[NEET-2013]

- (1) XeO_4 molecule is square planar.
- (2) There are four $p\pi-d\pi$ bonds.
- (3) There are four sp^3-p , σ bonds.
- (4) XeO_4 molecule is tetrahedral.

Q.18 Why do not helium, neon and argon form chemical compounds?

A.18 Since the ionization enthalpies of He, Ne and Ar are sufficiently higher than those of oxygen and fluorine.



Chemistry of Xenon


Xe reacts directly only with F_2 . Oxygen compounds can be obtained from the fluoride.

1. Xenon fluorides are white solid which are extremely powerful oxidising and fluorinating agents. They sublime readily at room temperature and are hydrolysed even by traces of water. Thus, they are stored in Ni containers.

Concept Ladder




Structures of Xenon fluorides cannot be explained by valence bond approach.






XeO₃ is a hygroscopic white explosive solid.
XeOF₄ is a colourless volatile liquid.



2. All xenon fluorides react with hydrogen and oxidise Cl^- to Cl_2 , I^- to I_2 and Ce(III) to Ce(IV)




3. Xenon fluorides reacts with fluoride ion acceptors to form cationic species and fluoride ion donors to form fluoro anions.




(M = Na, K, Rb, Cs)

4. Xe compounds show a variety of shapes:

XeF_2 is linear, XeF_4 is square planar, XeF_6 is a capped octahedron, XeO_3 is pyramidal, XeOF_4 is square pyramidal, XeO_2F_2 is see saw shaped, XeOF_2 is bent T - shaped, XeO_4 is tetrahedral, XeO_3F_3 is trigonal bipyramidal and $[\text{XeO}_6]^{4-}$ ion is octahedral.



XeF₂: Linear



XeF₄: Square planar



XeF₆: Distorted octahedral

Previous Year's Questions



Which compound has planar structure?

[AIPMT]

- (1) XeF_4 (2) XeOF_2
 (3) XeO_2F_2 (4) XeO_4

Q.19 What prompted Bartlett to the discovery of noble gas compounds?

A.19 Since PtF_6 oxidises O_2 to O_2^+ , Bartlett thought that PtF_6 should also oxidise Xe to Xe^+ because the ionization enthalpies of O_2 (1175 kJ mol^{-1}) and Xe (1170 kJ mol^{-1}) are quite close.

Q.20 In the preparation of H_2SO_4 by Contact Process, why is SO_3 not absorbed directly in water to form H_2SO_4 ?

A.20 A dense fog of H_2SO_4 is formed which is difficult to condense.

Q.21 PH_3 forms bubbles when passed slowly in water but NH_3 dissolves. Explain why?

A.21 Due to high electronegativity (3.0) and small size of N, NH_3 forms H-bonds with water and hence it is water soluble. Whereas, due to its lower electronegativity (2.1) of P and its bigger size than N, PH_3 does not form H-bonds with H_2O . As a result, it does not dissolve in H_2O and hence escapes as bubbles.

Q.22 In PCl_5 , phosphorus is in sp^3d hybridised state but all its five bonds are not equivalent. Justify your answer with reason.

A.22 Since each axial P—Cl bond is repelled by three bond pairs and each equatorial P—Cl bond is repelled by only two bond pairs, therefore, axial bonds are longer (240 pm) than equatorial bonds (202 pm). Thus, all the five P—Cl bonds in PCl_5 are not equivalent.

Q.23 Why is nitric oxide paramagnetic in gaseous state but the solid obtained on cooling it is diamagnetic?

A.23 Nitric acid has 15 electrons i.e. it has one unpaired electron and hence it is paramagnetic in the gaseous state. but in both solid as well as in liquid it exists as a dimer and hence is diamagnetic in nature.

Q.24 Give the reason to explain why ClF_3 exists but FCl_3 does not exist.

A.24 Because of bigger size, Cl can accommodate three small F atoms around it while F being smaller cannot accommodate three bigger sized Cl atoms around it.

Q.25 SF_6 is known but SH_6 is not. Why?

A.25 H_2 being a very weak oxidising agent cannot oxidise S to its maximum oxidation state of +6 but in case of fluorine (strongest oxidising agent) it oxidises S to its maximum oxidation state of +6, thus SF_6 is known but SH_6 is not known.

Q.26 The majority of noble gas compounds are those of xenon. Explain.

A.26 Except Rn which is radioactive, Xe has the lowest I.E. among inert gases and therefore it can be easily oxidised by strong oxidising agents like O_2 and F_2 . That is why majority of inert gas compounds are those of Xe.

Q.27 Arrange the hydrides of group 17 elements in order of their increasing acid strength.

A.27 As the size of halogen increases down the group from F to I, the bond dissociation energy of the H—X bond decreases. Consequently, the tendency of H—X bond to release a proton in water increases. In other words, the acid strength increases in the order : $HF < HCl < HBr < HI$.

Q.28 Why does argon not form diatomic molecules like oxygen and nitrogen?

A.28 All the orbitals (which are occupied by the electrons) are completely filled in Ar and hence it has no tendency to share electron with other atom of argon to form diatomic molecules. Oxygen and Nitrogen on the other hand, have unpaired electrons and hence share these electrons with other O and N atom respectively forming O_2 and N_2 molecules



Summary



Group 15 (N-family)

1. Hydrides

- (i) Bond angle : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$
 - (ii) Basic character : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$
 - (iii) Boiling point : $\text{PH}_3 > \text{AsH}_3 > \text{NH}_3 > \text{SbH}_3 > \text{BiH}_3$
 - (iv) Thermal stability : $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$
 - (v) Reducing character : $\text{NH}_3 < \text{PH}_3 < \text{AsH}_3 < \text{SbH}_3 < \text{BiH}_3$
2. (i) NF_3 is not hydrolysed due to absence of vacant d-orbital.
(ii) PF_3 and PF_5 are not hydrolysed because P—F bond is stronger than P—O bond.
 3. NF_3 is not lewis base although it has lone pair of e^- due to high electronegativity of fluorine.
 4. N_2O is called laughing gas.
 5. P_4S_3 is used in strike anywhere matches.
 6. Fuming HNO_3 is conc. $\text{HNO}_3 + \text{NO}_2$.
 7. PH_3 Poisonous gas having fish like smell.
 8. Holmes signal ships is $\text{CaC}_2 + \text{Ca}_3\text{P}_2$
 9. Match box side has red P + Sb_2S_3 + sand. Tip of match stick has $\text{K}_2\text{Cr}_2\text{O}_7 + \text{S} + \text{white P}$ or KClO_3 or red Pb.

Group 16 (O-family)

1. Acidic character

- (i) $\text{SO}_2 > \text{SeO}_2 > \text{TeO}_2 > \text{PoO}_2$
 - (ii) $\text{SO} < \text{SO}_2 < \text{SO}_3$
2. (i) Pyrosulphuric acid ($\text{H}_2\text{S}_2\text{O}_7$) has S—O—S linkage.
(ii) Marshall's acid has O—O linkage.
(iii) $\text{S}_2\text{O}_8^{2-}$ does not have S—S linkage.
 3. Catalyst used in manufacture of H_2SO_4 by Lead Chamber process is oxides of nitrogen and in contact process is V_2O_5 .

Group 17 (X-family)

1. Electronegativity : $\text{F} > \text{Cl} > \text{Br} > \text{I}$
2. Bond energy : $\text{Cl}_2 > \text{Br}_2 > \text{F}_2 > \text{I}_2$
3. Oxidising power : $\text{F}_2 > \text{Cl}_2 > \text{Br}_2 > \text{I}_2$
4. I_2 is soluble in water due to formation fo I_3^- polyhalide ion.

-
- Halide of metals in their higher oxidation states are more covalent than those formed in lower oxidation state. e.g. SnCl_4 is more covalent than SnCl_2 .
 - Hydrogen fluoride is not stored in glass bottles because it reacts with them. It is stored in copper, wax or guttapercha bottle.
 - KClO_3 is added in fire crackers as oxidising agent.
 - Tincher of iodine is $\text{I}_2 + \text{KI}$ dissolved in rectified spirit.
 - F_2 is called super halogen due to its high reactivity.
 - Br_2 or Cl_2 turn moist starch KI paper blue because they displace I_2 from KI and this I_2 reacts with starch to give blue complex.
 - AgF is soluble in water but AgCl , AgBr and AgI are not.

Group 18 (Noble gases)

- Discovery of noble gases.
He – Lockyer and Janassen
Ar – Lord Rayleigh and Ramsey
Kr – Ramsay and Travers
Xe – Ramsay and Travers
Ra – Doron
- Only He forms interstitial compounds with metals.
- O-He (1 : 4) mixture is used for treatment of asthma and for artificial respiration in deep sea diving because unlike N, He is not soluble in blood even at high pressure.